

HOMOGENEOUS IDEALS WITH MINIMAL SINGULARITY THRESHOLDS

BENJAMIN BAILY

ABSTRACT. Let $(\mathcal{O}_n, \mathfrak{m})$ denote the ring of germs of holomorphic functions $\mathbb{C}^n \rightarrow \mathbb{C}$, and let $I \subseteq \mathcal{O}_n$ be an \mathfrak{m} -primary ideal. Demailly and Pham showed that $\text{lct}(I) \geq \frac{1}{e_1(I)} + \cdots + \frac{e_{n-1}(I)}{e_n(I)}$, where $e_j(I)$ is the mixed multiplicity $e(I, \dots, I, \mathfrak{m}, \dots, \mathfrak{m})$, with I repeated j times and \mathfrak{m} repeated $n - j$ times.

We generalize the lower bound to the case of an arbitrary ideal of an excellent regular local (or standard-graded) ring of equal characteristic, with $\text{lct}(I)$ replaced by the F -threshold $c^{\text{m}}(I)$ in positive characteristic. Our main result is a classification of homogeneous ideals in polynomial rings for which the lower bound is attained, resolving a conjecture of Bivià-Ausina in the graded case.

1. INTRODUCTION

Let X be a smooth variety, $Y \subseteq X$ a proper closed subscheme, and $y \in Y$ a point. We study numerical invariants measuring the singularities of the pair (X, Y) at the point y : in characteristic zero, we consider the log canonical threshold; in positive characteristic, we consider the F -pure threshold.

The log canonical threshold (lct) has attracted considerable attention in algebraic geometry due to its connections with the Minimal Model Program and singularity theory. The F -pure threshold (fpt) is an analog of the lct in positive characteristic, related via reduction to characteristic $p \gg 0$ [HT04]. The fpt has provided insights into positive characteristic geometry [BS15; Kad+22; PRS24] and reduction modulo p has been leveraged to better understand singularities in characteristic zero [Ben+15; Tak04; Tak06].

In recent years, many authors have found lower bounds on the lct (and fpt) in terms of more classical invariants of the embedding [EM21; FEM03; FEM04; Kad+22; Kim21; Mus02; TW04]. The first of these bounds is due to Skoda [Sko72], who showed that

$$(1) \quad \frac{1}{\text{mult}_y(Y)} \leq \text{lct}_y(X, Y).$$

The case of equality in Equation (1) has long been known to experts: $\frac{1}{\text{mult}_y(Y)} = \text{lct}_y(X, Y)$ if and only if Y is a smooth divisor at y . Skoda's original formulation of Equation (1) concerned germs of plurisubharmonic functions, and the equality case

The author was supported by NSF grants DMS-2101075 and DMS-1840234 and a Simons Dissertation Fellowship.

in the analytic formulation was settled by Guan and Zhou [GZ15]. We consider the equality case in a generalization of Skoda's lower bound.

Let I denote the defining ideal of Y in $\mathcal{O}_{X,y}$ and $n = \dim \mathcal{O}_{X,y}$. In the case that $\mathcal{O}_{X,y}/I$ is zero-dimensional, Demailly and Pham [DP14] strengthened Equation (1) by incorporating higher-codimension information about the embedding, namely the mixed multiplicities $e_j(I) := e(I, \dots, I, \mathfrak{m}_y, \dots, \mathfrak{m}_y)$; see Definition 2.32. We have

$$(2) \quad \frac{1}{e_1(I)} + \frac{e_1(I)}{e_2(I)} + \dots + \frac{e_{n-1}(I)}{e_n(I)} \leq \text{lct}_y(X, Y).$$

Moving beyond the case considered by Demailly and Pham, we relax the assumption that $\mathcal{O}_{X,y}/I$ is zero-dimensional. We recall Rashkovskii's definition of the invariant $E_l(I)$ for any $l \leq \text{ht}(I)$. As in [Ras15, Theorem 1.4], we may let $L = V(h_{l+1}, \dots, h_n)$ denote the vanishing locus of $n - l$ general linear forms in $\mathcal{O}_{X,y}$ and let $I|_L$ denote the expansion of I to the ring $\mathcal{O}_{L,y}$. As $\mathcal{O}_{L,y}/I|_L$ is zero-dimensional, we apply Equation (2) and [Laz04, Example 9.5.4] to $I|_L$. This yields

$$(3) \quad \frac{1}{e_1(I|_L)} + \dots + \frac{e_{l-1}(I|_L)}{e_l(I|_L)} \leq \text{lct}(I|_L) \leq \text{lct}(I).$$

We then define

$$E_l(I) := \frac{1}{e_1(I|_L)} + \dots + \frac{e_{l-1}(I|_L)}{e_l(I|_L)},$$

which is well-defined for sufficiently general L by Proposition 2.37. The content of Equation (3) is that $E_l(I) \leq \text{lct}(I)$.

More generally, if (R, \mathfrak{m}) is an arbitrary regular local ring and $I \subseteq R$ with $\text{ht}(I) \geq l$, we define $E_l(I)$ similarly — in particular, when R/\mathfrak{m} is infinite, we let L denote the vanishing set of $n - l$ elements of \mathfrak{m} whose images in $\mathfrak{m}/\mathfrak{m}^2$ are sufficiently general and set $E_l(I) := E_l(I|_L)$. See Definition 2.38 for a precise definition.

Our first contribution is a generalization of Equation (3) to arbitrary characteristic.

Theorem A (Theorem 3.9). *Let (R, \mathfrak{m}) be a regular local ring or a polynomial ring over a perfect field considered with the standard grading. When $\text{char } R = 0$, assume that R is excellent. Let $I \subseteq R$ be an ideal of height at least l , which is assumed to be either homogeneous or \mathfrak{m} -primary when R is a polynomial ring. Let $c(I)$ denote the lct (resp. fpt) of I . Then $E_l(I) \leq c(I)$.*

In contrast to Skoda's bound, little is known about the case of equality in Equation (2). The best result in this direction is due to Bivià-Ausina [BA17], who classified cases of equality under the additional assumption that $\text{lct}(I) = \text{lct}(I^0)$, where I^0 is the smallest monomial ideal containing I . Our main result is coordinate-agnostic and resolves the conjecture of [BA24, Remark 3.7] in the graded case.

Theorem B (Theorem 5.1). *Let k be a perfect field of characteristic zero (resp. $p > 0$). Let $R = k[x_1, \dots, x_n]$ and let $I \subseteq R$ be a homogeneous ideal of height at least l . If $E_l(I) = \text{lct}(I)$ (resp. $E_l(I) = \text{fpt}(I)$), then there exist integers d_1, \dots, d_l*

such that, in suitable coordinates, we have

$$\bar{I} = \overline{(x_1^{d_1}, \dots, x_l^{d_l})}.$$

We briefly outline the proof of Theorem 5.1, which reduces to the case that $k = \bar{k}$ and $l = n$. Assume $E_n(I) = c(I)$. Write $I = I_1 + \dots + I_r$, where I_j is generated by d_j -forms and $d_1 < \dots < d_r$.

- (1) Using techniques from [BA17; May14], we control the generic initial ideals $\{\text{gin}(I^n)\}_{n \geq 1}$.
- (2) Using (1), we obtain a formula for the numbers $e_j(I)$ in terms of the d_j and the heights of the ideals $I_1 + \dots + I_j$.
- (3) Our formula for the $e_j(I)$ implies that $E_n(I) = B(I)$, where $B(I)$ is defined in terms of mixed Łojasiewicz exponents as in [BA24]. This allows us to reduce to the case of a homogeneous complete intersection.
- (4) We prove the complete intersection case (Proposition 4.30) by induction on the number of distinct degrees d_1, \dots, d_r .

In the case $r = 1$, any \mathfrak{m} -primary ideal I generated by d -forms automatically satisfies $\bar{I} = \mathfrak{m}^d$, so there is no way to use $r = 1$ as a useful base case for our induction. Instead, we use $r = 2$. In this case, we show (Lemma 4.8) that $c(I) = E_n(I)$ if and only if $c(I_1) = \text{ht}(I_1)/d_1$. As $\bar{I} = \overline{I_1 + \mathfrak{m}^{d_2}}$, Theorem B is equivalent in this case to the claim that $\bar{I}_1 = (x_1, \dots, x_{\text{ht}(I_1)})^{d_1}$ in suitable coordinates. In characteristic zero, this follows from [FEM03, Theorem 3.5]. In positive characteristic, this fact is [Bai25, Theorem 3.17].

By Example 5.14, the assumption that k is perfect cannot be weakened. The other main assumption in Theorem 5.1 is that R is graded and I is homogeneous, the relaxation of which is a subtler question. In characteristic zero, we conjecture that Theorem 5.1 holds in the local case up to an *analytic* change of coordinates; see Conjecture 6.2 for the precise statement and Conjecture 6.3 for an even stronger statement in terms of valuations. We verify both conjectures in dimension 2. In contrast, in characteristic $p > 0$, Example 6.1 shows that analytic changes of coordinates do not suffice to characterize when $c(I) = E_l(I)$.

We also remark on two (a priori) possible improvements to Theorems A and B. The first is regarding [Hie25, Theorem 1.2], which defines for each $1 \leq k \leq n$ an invariant $H_k(I)$ satisfying $E_k(I) \leq H_k(I) \leq \text{lct}(I)$ for all ideals $I \subseteq \mathcal{O}_n$ of height at least k . It would be interesting to see whether the bound $H_k(I) \leq c(I)$ remains true in positive characteristic, and moreover whether there exist ideals for which $E_k(I) < H_k(I) = c(I)$.

The second remark concerns [Kim21, Question 1.3], which conveys a conjecture of Hoang Hiep Pham that (implies) $c(I) - c(I|_H) \geq \frac{e_{n-1}(I)}{e_n(I)}$ for every height- n ideal $I \subseteq \mathcal{O}_n$ and every hyperplane H through $0 \in \mathbb{C}^n$. Tommaso de Fernex shared a counterexample to this conjecture with the author, which we record in Example 3.13.

Acknowledgments. The author is grateful to his advisor, Karen Smith, for her support and guidance through many iterations of this project. The author is additionally grateful to Havi Ellers and Mattias Jonsson for feedback on an earlier draft, and to Tommaso de Fernex for helpful discussions.

2. PRELIMINARIES

Assumption 2.1. *Throughout this article, all rings are commutative, Noetherian, and unital. All polynomial rings are considered with the standard grading.*

2.1. F -Pure and Log Canonical Thresholds. We begin with a formal definition of the lct. For a detailed introduction, see [Mus12].

Definition 2.2 (Log Resolution). Let X be a smooth variety and $Y \subseteq X$ a proper closed subvariety with defining ideal \mathfrak{a} . Let W be a smooth variety. A projective morphism $\pi : W \rightarrow X$ is a **log resolution** of (X, Y) if π is an isomorphism over $X \setminus Y$ and the inverse image $\mathfrak{a} \cdot \mathcal{O}_W$ is the ideal of a Cartier divisor D such that $D + K_{W/X}$ has simple normal crossings.

Definition-Proposition 2.3 (Log canonical threshold, [Mus12] Theorem 1.1). Let X be a smooth variety over a characteristic zero field and $Y \subseteq X$ a closed subvariety with defining ideal \mathfrak{a} . By Hironaka's theorem on resolution of singularities, there exists a log resolution $\pi : W \rightarrow X$ of the pair (X, Y) . In particular, there exists a simple normal crossings divisor $E_1 + \cdots + E_N$ such that

$$D = \sum_{i=1}^N a_i E_i \quad \text{and} \quad K_{W/X} = \sum_{i=1}^N k_i E_i.$$

For $y \in Y$, the quantity $\min_{i: y \in \pi(E_i)} \frac{k_i + 1}{a_i}$ does not depend on the log resolution π and is denoted $\text{lct}_y(X, Y)$, the **log canonical threshold** of (X, Y) at y . If a point y is not specified, we define

$$\text{lct}(X, Y) = \min_{y \in Y} \text{lct}_y(X, Y) = \min_i \frac{k_i + 1}{a_i}.$$

We make the convention that $\text{lct}_y(X, Y) = \infty$ when $y \notin Y$. Additionally, when k is a characteristic zero field, $R = k[x_1, \dots, x_n]$, and $I \subseteq R$ is an ideal, we set $\text{lct}(R, I) := \text{lct}(\mathbb{A}^n, V(I))$ and $\text{lct}_0(R, I) := \text{lct}_0(\mathbb{A}^n, V(I))$.

In fact, one can define the log canonical threshold of (X, Y) whenever X is an integral *excellent* scheme; see [FM09, §2].

For background on the F -pure threshold, we direct the reader to [TW04]. In this subsection, we summarize several key definitions and results.

Definition 2.4. Let R be a ring of characteristic $p > 0$. We let F_*R denote the R -module structure on R given by restriction of scalars along the Frobenius map $F : R \rightarrow R$. We say R is F -finite if F_*R is module-finite over R .

Definition 2.5 ([TW04]). Let R be an F -finite ring, $I \subseteq R$ an ideal, and $t \in \mathbb{R}_{\geq 0}$. The pair (R, I^t) is F -split if for $e \gg 0$, the map

$$I^{\lfloor t(p^e-1) \rfloor} \cdot \text{Hom}(F_*^e R, R) \rightarrow R$$

is surjective. The F -pure threshold of the pair (R, I) is the supremum of all t such that (R, I^t) is F -split. We denote this quantity by $\text{fpt}(R, I)$, or $\text{fpt}(I)$ when the ambient ring is clear.

Let k be an F -finite field of characteristic $p > 0$, $R = k[x_1, \dots, x_n]$, and $I \subseteq R$. Let $\mathfrak{m} := (x_1, \dots, x_n)$ denote the homogeneous maximal ideal of R . We then let $\text{fpt}_0(I)$ denote the quantity $\text{fpt}(R_{\mathfrak{m}}, IR_{\mathfrak{m}})$.

In practice, we do not use the above definitions. Instead, we use the characterization of the F -pure threshold as the F -threshold at the maximal ideal and the lct as a limit of fpt .

Definition 2.6. Let R be a ring of characteristic $p > 0$. Let \mathfrak{a}, J be ideals of R such that $\mathfrak{a} \subseteq \sqrt{J}$. For each nonnegative integer e , we define

$$\nu_{\mathfrak{a}}^J(p^e) := \max\{t \in \mathbb{Z}^+ : \mathfrak{a}^t \not\subseteq J^{\lfloor p^e \rfloor}\}.$$

By [DSBP18], the sequence $\frac{\nu_{\mathfrak{a}}^J(p^e)}{p^e}$ has a limit as $e \rightarrow \infty$; we denote this limit by $c^J(\mathfrak{a})$ and refer to it as the F -threshold of \mathfrak{a} at J .

Proposition 2.7. *Let (R, \mathfrak{m}) be an F -finite regular local ring. Then the F -pure threshold of the pair (R, I) is equal to $c^{\mathfrak{m}}(I)$. If instead R is a polynomial ring over an F -finite field and $I \subseteq R$ is a homogeneous ideal, then the same holds when we let \mathfrak{m} denote the homogeneous maximal ideal of R .*

Proof. The local case is [MTW05, Remark 1.5] and the graded case is [DSB18, Proposition 3.10]. \square

Proposition 2.8 ([HY03], Theorem 6.8). *Let A be a finite-type \mathbb{Z} -algebra and $\mathfrak{a} \subseteq A[x_1, \dots, x_n]$ an ideal. Set $k = \text{Frac}(A)$. Then we have*

$$\text{lct}(\mathfrak{a} \otimes_A k) = \lim_{\mu \in \max \text{Spec } A, |A/\mu| \rightarrow \infty} \text{fpt}(A/\mu[x_1, \dots, x_n], \mathfrak{a} \otimes_A A/\mu).$$

Many of our results make sense for both fpt and lct , so we introduce the following notation to avoid stating the same results once each for characteristic zero and positive characteristic.

Notation 2.9. *Let k be a field, $R = k[x_1, \dots, x_n]$, $\mathfrak{m} = (x_1, \dots, x_n)$, and $I \subseteq R$ an ideal. We define the quantity $c(R, I)$ as follows:*

$$c(R, I) = \begin{cases} c^{\mathfrak{m}}(I) & \text{char } R = p > 0 \\ \text{lct}_0(R, I) & \text{char } R = 0. \end{cases}$$

Similarly, let (R, \mathfrak{m}) be a regular local ring and $I \subseteq R$ an ideal. We define

$$c(R, I) = \begin{cases} c^{\mathfrak{m}}(I) & \text{char } R = p > 0 \\ \text{lct}(R, I) & \text{char } R = 0 \text{ and } R \text{ is excellent.} \end{cases}$$

If the ambient ring is clear, we will use $c(I)$ as shorthand.

Proposition 2.10 (Properties of the singularity threshold). *Assume either setting of Notation 2.9, and let I, J be nonzero ideals of R . The following properties hold for $c(-)$:*

- (i) *If $I \subseteq J$, then $c(I) \leq c(J)$.*
- (ii) *For all $m > 0$, we have $c(I^m) = m^{-1}c(I)$.*
- (iii) *We have $c(I) = c(\bar{I})$, where \bar{I} denotes the integral closure of I .*
- (iv) *If $J \subseteq R$ is an ideal such that R/J is regular, then*

$$c(R, I) \geq c(R/J, (I + J)/J).$$

- (v) *We have $c(I + J) \leq c(I) + c(J)$.*
- (vi) *Suppose $R \subseteq S$ is a ring extension. If $\text{char } R = 0$, assume S is excellent. In both of the following cases, $c(I) = c(IS)$.*
 - *(R, \mathfrak{m}) and (S, \mathfrak{n}) are local and $\mathfrak{m}S = \mathfrak{n}$;*
 - *$R = k[x_1, \dots, x_n]$, L/k is a field extension, and $S = L[x_1, \dots, x_n]$.*

Proof. First, we verify (i)-(v). For characteristic zero, see [Mus12, Properties 1.12, 1.13, 1.15, 1.17, 1.20]. For characteristic $p > 0$, see [TW04, Proposition 2.2 (1), (2), (6), Proposition 4.4].

For (vi), note that both cases are regular and faithfully flat extensions. In characteristic zero, see [JM12, Proposition 1.9]. In characteristic $p > 0$, see [Hun+08, Proposition 2.2 (v)]. \square

2.2. Monomial Ideals.

Notation 2.11. *For a vector of ring elements $\mathbf{f} = f_1, \dots, f_r$ and a vector of non-negative integers $\mathbf{a} = a_1, \dots, a_r$, we let $\mathbf{f}^{\mathbf{a}}$ denote the element $f_1^{a_1} \dots f_r^{a_r}$. A boldface, lowercase letter always refers to a vector of integers or ring elements.*

Definition 2.12. Let k be a field and let $R = k[x_1, \dots, x_n]$. For a tuple $\mathbf{x} = x_{i_1}, \dots, x_{i_r}$ with $1 \leq i_1 < \dots < i_r \leq n$, we let $\text{Mon}(\mathbf{x})$ denote the monoid of monomials in the variables \mathbf{x} . In particular, $\text{Mon}(x_1, \dots, x_n)$ denotes the set of all monomials in R .

When working with monomial ideals, one often identifies a monomial $x_1^{a_1} \dots x_n^{a_n}$ with the point $(a_1, \dots, a_n) \in \mathbb{Z}_{\geq 0}^n$. For future reference, it will help to give a name to this identification.

Definition 2.13. Let k be a field. We define the map

$$\log : \text{Mon}(x_{i_1}, \dots, x_{i_r}) \rightarrow \mathbb{Z}_{\geq 0}^r, \quad \log(x_{i_1}^{a_1} \dots x_{i_r}^{a_r}) = (a_1, \dots, a_r).$$

Definition 2.14. Let $\mathfrak{a} \subseteq k[x_1, \dots, x_n]$ be a monomial ideal. Then the *Newton Polytope* of \mathfrak{a} , denoted $\Gamma(\mathfrak{a})$, is the convex hull in $\mathbb{R}_{\geq 0}^n$ of $\log(\mathfrak{a})$. We let $\text{conv}(-)$ denote the convex hull of a set.

Remark 2.15. We record several properties of $\Gamma(\mathfrak{a})$.

- (i) $\Gamma(\mathfrak{a})$ is a closed, convex, unbounded subset of the first orthant of \mathbb{R}^n .




FIGURE 1. Computation of μ for Example 2.17

- (ii) When \mathfrak{a} is an \mathfrak{m} -primary ideal, the closure of the complement of $\Gamma(\mathfrak{a})$ inside the first orthant is an open, bounded polyhedron.
- (iii) For two ideals $\mathfrak{a}, \mathfrak{b}$, the Minkowski sum of $\Gamma(\mathfrak{a})$ and $\Gamma(\mathfrak{b})$ is equal to $\Gamma(\mathfrak{a}\mathfrak{b})$. In particular, $\Gamma(\mathfrak{a}^n) = n\Gamma(\mathfrak{a})$.

The following proposition shows that Newton polytope of a monomial ideal determines the singularity threshold.

Proposition 2.16. *Let $\mathfrak{a} \subseteq k[x_1, \dots, x_n]$ be a monomial ideal. Then*

$$c(\mathfrak{a}) = \frac{1}{\mu}, \text{ where } \mu = \inf\{t : t\mathbf{1} \in \Gamma(\mathfrak{a})\}.$$

Proof. See [How01], Example 5 for characteristic zero and [Her16], Proposition 36 for positive characteristic. \square

Example 2.17. Let $R = k[x, y]$ and set $\mathfrak{a} = (x^6, x^5y, x^3y^2, x^2y^3, xy^4, y^6)$. In Figure 1 we compute $\mu = 2.5$, so $c(\mathfrak{a}) = 0.4$.

Following the proof of [FEM04], Theorem 1.4 and the terminology of [May14], we also define the **limiting polytope** of a graded system of monomial ideals.

Definition 2.18. Let \mathfrak{a}_\bullet be a graded system of monomial ideals in $k[x_1, \dots, x_n]$ — that is, suppose $\mathfrak{a}_r\mathfrak{a}_s \subseteq \mathfrak{a}_{r+s}$ for all $r, s \in \mathbb{Z}^+$. We define $\Gamma(\mathfrak{a}_\bullet)$ as the closure in \mathbb{R}^n of the ascending union $\{\frac{1}{2^m}\Gamma(\mathfrak{a}_{2^m})\}_{m>0}$.

Remark 2.19. The set $\overline{\bigcup_m \frac{1}{2^m}\Gamma(\mathfrak{a}_{2^m})}$ in Definition 2.18 coincides with $\overline{\bigcup_m \frac{1}{m}\Gamma(\mathfrak{a}_m)}$ in the settings we consider (where $\mathfrak{a}_m \supseteq \mathfrak{a}_{m+1}$), but we find it easier to work with an ascending union than a filtered union.

We fix our conventions for working with monomial orders and initial terms in polynomial rings. For further background, see [Eis95, Chapter 15].

Definition 2.20. Let k be a field and let $R = k[x_1, \dots, x_n]$. A *monomial order* $>$ on R is a partial order on $\text{Mon}(x_1, \dots, x_n)$ compatible with the divisibility relations: that is, for monomials $\mathbf{x}^{\mathbf{a}}, \mathbf{x}^{\mathbf{b}}, \mathbf{x}^{\mathbf{c}}$, if $\mathbf{x}^{\mathbf{a}} > \mathbf{x}^{\mathbf{b}}$ then $\mathbf{x}^{\mathbf{a}+\mathbf{c}} > \mathbf{x}^{\mathbf{b}+\mathbf{c}}$.

For an element $f \in R$, write $f = \sum_{\mathbf{a} \in \mathbb{Z}_{\geq 0}^n} \beta_{\mathbf{a}} \mathbf{x}^{\mathbf{a}}$ where $\beta_{\mathbf{a}} \in k$. The **support** of f , denoted $\text{supp}(f)$, is the set $\{\mathbf{x}^{\mathbf{a}} : \beta_{\mathbf{a}} \neq 0\}$. The *initial term* of f with respect to $>$, denoted $\text{in}_{>}(f)$, is given by

$$\sum_{\substack{\mathbf{a}: \mathbf{x}^{\mathbf{a}} \in \text{supp}(f) \\ \mathbf{x}^{\mathbf{a}} \text{ maximal with respect to } >}} \beta_{\mathbf{a}} \mathbf{x}^{\mathbf{a}}.$$

For an ideal $I \subseteq R$, the *initial ideal* $\text{in}_{>}(I)$ is the ideal generated by the elements $\{\text{in}_{>}(f) : f \in I\}$.

Valuations provide an important source of monomial orders.

Definition 2.21. Assume the setting of Definition 2.20. A *monomial valuation* is a map of monoids $v : (\text{Mon}(x_1, \dots, x_n), \cdot) \rightarrow (\mathbb{R}, +)$. To describe a monomial valuation v , it suffices to give the values $v(x_1), \dots, v(x_n)$. We can then describe a monomial order $>_v$ on R by

$$\mathbf{x}^{\mathbf{a}} >_v \mathbf{x}^{\mathbf{b}} \text{ if and only if } v(\mathbf{x}^{\mathbf{a}}) > v(\mathbf{x}^{\mathbf{b}})$$

When $v(x_i) \geq 0$ for all i , we can extend v to an \mathbb{R} -valuation on R : we define $v : R \rightarrow [0, \infty]$ by

$$v(f) = \min_{\mathbf{x}^{\mathbf{a}} \in \text{supp}(f)} v(\mathbf{x}^{\mathbf{a}}).$$

A crucial property of initial ideals is semicontinuity with respect to the singularity threshold.

Proposition 2.22. Let $R = k[x_1, \dots, x_n]$, $\mathfrak{m} = (x_1, \dots, x_n)$ and let $>$ be a monomial order. If $I \subseteq R$ is an ideal, then $c(\text{in}_{>}(I)) \leq c(I)$.

Proof. For characteristic zero, see [DK01] for the semicontinuity of the let. For positive characteristic, see the claim preceding [TW04, Remark 4.6]. \square

When I is homogeneous, k is infinite, and $>$ is a total order compatible with the partial order by degree, there is a well-defined notion of the initial ideal in “generic coordinates.”

Definition-Proposition 2.23 ([Eis95], Theorem 15.18). Let k be an infinite field, $R = k[x_1, \dots, x_n]$, and $I \subseteq R$ a homogeneous ideal. Suppose that $>$ is a monomial total order on R such that $x_1 > \dots > x_n$ and $\mathbf{x}^{\mathbf{a}} > \mathbf{x}^{\mathbf{b}}$ whenever $\deg(\mathbf{x}^{\mathbf{a}}) > \deg(\mathbf{x}^{\mathbf{b}})$. Then there exists a nonempty open subset $U \subseteq \text{GL}_n(k)$ and a monomial ideal J such that:

- For all $\gamma \in U$, we have $\text{in}_{>}(\gamma I) = J$;
- U is *Borel-fixed*: if $B \subseteq \text{GL}_n(k)$ denotes the subgroup of lower triangular matrices, then $BU = U$.

The ideal J is called the *generic initial ideal* of I with respect to $>$, and is denoted $\text{gin}_{>}(I)$.

2.3. Integral Closure of Ideals.

Definition 2.24. Let I be an ideal in a ring R . An element $r \in R$ is integral over I if there exists an integer n and elements $a_1, \dots, a_n, a_i \in I^i$ such that

$$r^n + a_1 r^{n-1} + \dots + a_n = 0.$$

We then define the integral closure \bar{I} of I as the set of elements $r \in R$ which are integral over I .

Definition-Proposition 2.25 ([HS06], Corollary 1.2.5). Let $J \subseteq I$ be ideals in ring R . We say J is a **reduction** of I if either of the following equivalent conditions holds:

- (i) There exists a positive integer n such that $I^n J = I^{n+1}$.
- (ii) We have $I \subseteq \bar{J}$;
- (iii) We have $\bar{I} = \bar{J}$.

Those hoping for an exhaustive discussion of the integral closure of ideals should consult [HS06]. For now, we list some basic properties of \bar{I} .

Proposition 2.26 (Properties of the Integral Closure, [HS06] Chapter 1). *Let R be a ring and $I \subseteq R$ an ideal. Let $\varphi : R \rightarrow S$. Then we have*

- (i): \bar{I} is an ideal.
- (ii): $\overline{(\bar{I})} = \bar{I}$.
- (iii): $\bar{I}S \subseteq \overline{IS}$.
- (iv): If $J \subseteq S$ is an ideal, then $\varphi^{-1}(\bar{J}) = \overline{\varphi^{-1}(J)}$.
- (v): For any multiplicatively closed subset $W \subseteq R$, we have $W^{-1}\bar{I} = \overline{W^{-1}I}$.
- (vi): The integral closure of a monomial ideal \mathfrak{a} in a polynomial ring $k[x_1, \dots, x_n]$ is generated by the set $\mathbf{x}^{\mathbf{a}} : \mathbf{a} \in \Gamma(\mathfrak{a})$.
- (vii): If φ is faithfully flat or an integral extension, then $\bar{I}S \cap R = \bar{I}$.

Integral closure is an operation which respects many numerical invariants we are interested in this paper.

Theorem 2.27 ([HS06], Proposition 11.2.1, Theorem 11.3.1). *Let (R, \mathfrak{m}) be a formally equidimensional local ring and $I \subseteq J$ two \mathfrak{m} -primary ideals. Then $e(I) = e(J)$ if and only if $\bar{I} = \bar{J}$.*

The same result holds in the case that (R, \mathfrak{m}) is instead standard-graded.

Proposition 2.28. *Let $I \subseteq k[x_1, \dots, x_n]$ be an ideal. Then $c(I) = c(\bar{I})$.*

Proof. For characteristic zero, see [Mus12], Property 1.15. For positive characteristic, see [TW04], Proposition 2.2 (6). \square

To conclude this subsection, we recall a version of the Briançon-Skoda theorem, due to Lipman and Sathaye in the level of generality we need.

Theorem 2.29. *Let (R, \mathfrak{m}) be a regular local ring of dimension d and $I \subseteq R$ an ideal. Then for all integers $t > 0$, we have*

$$\bar{I}^{t+d} \subseteq \overline{I^{t+d}} \subseteq I^{t+1}.$$

Proof. The first containment is [HS06, Proposition 5.3.1]. For the second, we reduce to the case of an infinite residue field; we let $R(X) := R[X]_{\mathfrak{m}R[X]}$ as in op. cit. By Proposition 2.26 we have $\bar{I} = \overline{IR(X)} \cap R$, so it suffices to prove the claim for $R = R(X)$.

By [HS06, Proposition 5.1.6 and Theorem 8.6.6] there exists an ideal $J \subseteq I$ with $\overline{J^t} = \bar{I}^t$ for all $t > 0$ and such that J can be generated by at most d elements. By [LS81, Theorem 1''] we deduce

$$\overline{I^{t+d}} = \overline{J^{t+d}} \subseteq J^{t+1} \subseteq I^{t+1}.$$

□

2.4. Mixed Multiplicities and the Demailly-Pham Invariant. We recall the definition of the mixed multiplicity symbol $e(I_1, \dots, I_d; M)$.

Definition 2.30. Let M be a finite-length R -module. We let $\lambda_R(M)$ denote the length of M as an R -module.

Theorem 2.31 ([HS06], Theorem 17.4.2). *Let (R, \mathfrak{m}) be a Noetherian local ring, I_1, \dots, I_k ideals of R primary to \mathfrak{m} , and M a finitely-generated R -module. Then there exists a polynomial $P(n_1, \dots, n_k)$ with rational coefficients and total degree at most $\dim R$ such that for all $n_1, \dots, n_k \gg 0$, we have*

$$(4) \quad P(n_1, \dots, n_k) = \lambda_R \left(\frac{M}{I_1^{n_1} \dots I_k^{n_k} M} \right).$$

Definition 2.32 (Mixed Multiplicity). Let (R, \mathfrak{m}) be a Noetherian local ring of dimension d . Let I_1, \dots, I_k be \mathfrak{m} -primary ideals of R and let $P(n_1, \dots, n_k)$ be as in Equation (4). Write $Q(n_1, \dots, n_k)$ for the degree- d part of $P(n_1, \dots, n_k)$. The coefficients of Q define the **mixed multiplicities** $e(I_1^{(d_1)}, \dots, I_k^{(d_k)}; M)$:

$$(5) \quad Q(n_1, \dots, n_k) = \sum_{d_1 + \dots + d_k = d} \binom{d}{d_1, \dots, d_k}^{-1} e(I_1^{(d_1)}, \dots, I_k^{(d_k)}; M)$$

The expression $e(I_1^{(d_1)}, \dots, I_k^{(d_k)}; M)$ is shorthand for the expression

$$e(I_1, \dots, I_1, \dots, I_k, \dots, I_k, \dots, I_d; M),$$

where I_j is repeated d_j times.

Remark 2.33. Other authors [HS06] have used the notation $e(I_1^{[d_1]}, \dots, I_k^{[d_k]}; M)$. To avoid confusion with the Frobenius powers $I_j^{[p^e]}$ of the ideals I_j , we use angle brackets in the exponent.

We now define the mixed multiplicities $e_j(I)$.

Definition 2.34. Let (R, \mathfrak{m}) be a Noetherian local ring of dimension d and let I denote an \mathfrak{m} -primary ideal. We define

$$e_j(I) = e(I^{(j)}, \mathfrak{m}^{(d-j)}; R).$$

We record a few basic properties of the numbers $e_j(I)$.

Proposition 2.35. *Let (R, \mathfrak{m}) be a regular local ring of dimension n with infinite residue field. Let $I \subseteq R$ be an \mathfrak{m} -primary ideal.*

- (i) *We have $e_0(I) = 1$, $e_1(I) = \text{ord}_{\mathfrak{m}}(I)$, and $e_n(I) = e(I)$.*
- (ii) *If h_1, \dots, h_n are elements of \mathfrak{m} whose images in $\mathfrak{m}/\mathfrak{m}^2$ are sufficiently general, then for all $0 \leq j \leq n$ we have*

$$e_j(I) = e\left(\frac{I + (h_1, \dots, h_{n-j})}{(h_1, \dots, h_{n-j})}\right) = e(I + (h_1, \dots, h_{n-j})),$$

where $e(-)$ denotes the usual Hilbert-Samuel multiplicity.

- (iii) *Minkowski inequality: for $j \geq 1$, we have $e_j(I)^2 \leq e_{j-1}(I)e_{j+1}(I)$.*

Proof.

- (i) Follows from (ii).
- (ii) The first equality is [HS06, Corollary 17.4.7]; the second is [HS06, Proposition 11.1.19].
- (iii) Follows from [HS06, Theorem 17.7.2].

□

To state Theorem 3.9, we must extend Definition 2.32 to the case of an ideal $I \subseteq R$ which is not necessarily \mathfrak{m} -primary.

Definition 2.36 ([BA08], Definition 2.4). Let (R, \mathfrak{m}) be an n -dimensional local ring and I_1, \dots, I_n ideals of R . We define

$$(6) \quad \sigma(I_1, \dots, I_n) = \sup_{t>0} e(I_1 + \mathfrak{m}^t, \dots, I_n + \mathfrak{m}^t).$$

As a special case, for $1 \leq j \leq n$ and $I \subseteq R$, we define $\sigma_j(I) := \sigma(I^{(j)}, \mathfrak{m}^{\langle n-j \rangle})$ as in Definition 2.34.

The quantity (6) may be infinite. The following proposition summarizes basic properties of $\sigma(I_1, \dots, I_n)$, among which is a characterization of when the quantity (6) is finite.

Proposition 2.37 ([BA08]). *Let (R, \mathfrak{m}) be an n -dimensional regular local ring and I_1, \dots, I_n ideals of R .*

- (i) *If I_1, \dots, I_n have height n , then $\sigma(I_1, \dots, I_n) = e(I_1, \dots, I_n)$.*
- (ii) *We have $\sigma(I_1, \dots, I_n) < \infty$ if and only if there exist $g_1 \in I_1, \dots, g_n \in I_n$ such that (g_1, \dots, g_n) is \mathfrak{m} -primary. In this case, $\sigma(I_1, \dots, I_n) = e(g_1, \dots, g_n)$ for elements $g_i \in I_i$ whose images in $I_i/\mathfrak{m}I_i$ are sufficiently general.*
- (iii) *In particular, for an ideal $I \subseteq R$, we have $\sigma_j(I) < \infty$ if and only if $\text{ht}(I) \geq j$. In this case, if $h_1, \dots, h_{n-j} \in \mathfrak{m}$ are elements whose images in $\mathfrak{m}/\mathfrak{m}^2$ are sufficiently general, we have*

$$(7) \quad \sigma_j(I) = e\left(\frac{I + (h_1, \dots, h_{n-j})}{(h_1, \dots, h_{n-j})}\right) = e(I + (h_1, \dots, h_{n-j})).$$

- (iv) *The Minkowski inequalities hold: if $\sigma_{j-1}(I), \sigma_j(I)$ are finite then $\sigma_j(I)^2 \leq \sigma_{j-1}(I)\sigma_{j+1}(I)$.*

Proof.

- (i) For $t \gg 0$, we have $\mathfrak{m}^t \subseteq I_i$ for all $1 \leq i \leq n$. See the remark after Definition 2.4 in op. cit.
- (ii) This is Proposition 2.4 in op. cit.
- (iii) When $j \leq \text{ht } I$, use prime avoidance to choose $g_1, \dots, g_j \in I$ such that $\text{ht}((g_1, \dots, g_j)) = j$ and $g_{j+1}, \dots, g_n \in \mathfrak{m}$ such that $\text{ht}((g_1, \dots, g_n)) = n$. When $j \geq \text{ht}(I) + 1$, for any $g_1, \dots, g_j \in I, g_{j+1}, \dots, g_n \in \mathfrak{m}$ we have

$$\text{ht}((g_1, \dots, g_n)) \leq \text{ht}(I + (g_{j+1}, \dots, g_n)) \leq \text{ht}(I) + n - j \leq n - 1.$$

To see that the first equality in Equation (7) holds, pick $t \gg 0$ such that $\sigma_j(I) = e_j(I + \mathfrak{m}^t)$. Use (ii) to pick $g_1, \dots, g_j \in I, h_1, \dots, h_{n-j} \in \mathfrak{m}$ generally such that $\sigma_j(I) = e((g_1, \dots, g_j, h_1, \dots, h_{n-j}))$. Use Proposition 2.35(ii) to pick the h_1, \dots, h_{n-j} even more generally so that $e_j(I + \mathfrak{m}^t) = e(I + \mathfrak{m}^t + (h_1, \dots, h_{n-j}))$. Using monotonicity of Hilbert-Samuel multiplicity, we compute

$$\begin{aligned} \sigma_j(I) &= e_j(I + \mathfrak{m}^t) = e(I + \mathfrak{m}^t + (h_1, \dots, h_{n-j})) \leq e(I + (h_1, \dots, h_{n-j})) \\ &\leq e(g_1, \dots, g_j, h_1, \dots, h_{n-j}) = \sigma_j(I). \end{aligned}$$

- (iv) Follows immediately from Proposition 2.35 (iii). □

We will now define the Demailly-Pham invariant [DP14].

Definition 2.38. Let (R, \mathfrak{m}) be a regular local ring, $l \in \mathbb{Z}^+$, and I an ideal with $\text{ht}(I) \geq l$. Then we set

$$E_l(I) := \frac{1}{\sigma_1(I)} + \dots + \frac{\sigma_{l-1}(I)}{\sigma_l(I)}.$$

Suppose instead $R = k[x_1, \dots, x_n]$ and $\mathfrak{m} = (x_1, \dots, x_n)$. Let $I \subseteq R$ be an \mathfrak{m} -primary ideal. For all $r, s > 0$, we have

$$\lambda_R \left(\frac{R}{I^r \mathfrak{m}^s} \right) = \lambda_{R_{\mathfrak{m}}} \left(\frac{R_{\mathfrak{m}}}{I^r \mathfrak{m}^s R_{\mathfrak{m}}} \right),$$

so Theorem 2.31 holds for $\lambda_R \left(\frac{R}{I^r \mathfrak{m}^s} \right)$ without assuming that R is local. In particular, for $0 \leq j \leq n$, the numbers $e(I^{(j)}, \mathfrak{m}^{(n-j)})$ can be defined intrinsically from the polynomial $P(r, s)$ and agree with the quantities $e_j(IR_{\mathfrak{m}})$. Additionally, in this setting we have an analog of Theorem 2.27.

Proposition 2.39. *Let R be a regular ring and $\mathfrak{m} \subseteq R$ a maximal ideal such that $\dim R_{\mathfrak{m}} = n$. Let $I_1 \subseteq I_2 \subseteq R$ be \mathfrak{m} -primary ideals. Then $E_n(I_1 R_{\mathfrak{m}}) \leq E_n(I_2 R_{\mathfrak{m}})$ with equality if and only if $\overline{I_1} = \overline{I_2}$.*

Proof. The case that (R, \mathfrak{m}) is local is [BA17, Corollary 11]. In the non-local case, we have $\overline{I_1 R_{\mathfrak{m}}} = \overline{I_2 R_{\mathfrak{m}}}$, so $I_1 R_{\mathfrak{m}}$ is a reduction of $I_2 R_{\mathfrak{m}}$ by [HS06, Corollary 1.2.5].

By definition of reduction, there exists $t > 0$ such that $I_1 I_2^t R_{\mathfrak{m}} = I_2^{t+1} R_{\mathfrak{m}}$. The ideals $I_1 I_2^t, I_2^{t+1}$ are \mathfrak{m} -primary, so we have

$$I_1 I_2^t = I_1 I_2^t R_{\mathfrak{m}} \cap R = I_2^{t+1} R_{\mathfrak{m}} \cap R = I_2^{t+1},$$

so I_1 is a reduction of I_2 . By [HS06, Corollary 1.2.5] again, we deduce that $\overline{I_1} = \overline{I_2}$. \square

We extend Definition 2.38 to the ring $k[x_1, \dots, x_n]$.

Definition 2.40. Let $R = k[x_1, \dots, x_n]$, $\mathfrak{m} = (x_1, \dots, x_n)$ and let $I \subseteq \mathfrak{m}$ be an ideal. For $1 \leq j \leq \text{ht}(IR_{\mathfrak{m}})$ we define $\sigma_j(I) := \sigma_j(IR_{\mathfrak{m}})$ and $E_j(I) := E_j(IR_{\mathfrak{m}})$.

Remark 2.41. Even though Definition 2.40 defines a local invariant, we suppress the reference to the maximal ideal \mathfrak{m} , as we will only ever consider singularities at the origin.

Proposition 2.42. *Let $R = k[x_1, \dots, x_n]$, $\mathfrak{m} = (x_1, \dots, x_n)$, and let $I \subseteq R$ be an ideal with $\text{ht}(I) \geq l$.*

- (a) *If I is \mathfrak{m} -primary, then Proposition 2.35 holds for I .*
- (b) *If I is homogeneous, then Proposition 2.37 (i), (iii), (iv) hold, provided we assume the elements h_1, \dots, h_{n-j} in (iii) are homogeneous.*

Proof.

- (a) As in Proposition 2.35, (i) follows from (ii). For (ii), for any $h_1, \dots, h_{n-j} \in \mathfrak{m}$, the ideal $(I + (h_1, \dots, h_{n-j}))$ is \mathfrak{m} -primary as well. As $\mathfrak{m}/\mathfrak{m}^2 \cong \mathfrak{m}R_{\mathfrak{m}}/\mathfrak{m}^2R_{\mathfrak{m}}$, choosing $h_1, \dots, h_{n-j} \in \mathfrak{m}$ whose images in $\mathfrak{m}R_{\mathfrak{m}}/\mathfrak{m}^2R_{\mathfrak{m}}$ are sufficiently general, we have

$$e_j(I) := e_j(IR_{\mathfrak{m}}) = e(I + (h_1, \dots, h_{n-j})R_{\mathfrak{m}}) = e(I + (h_1, \dots, h_{n-j})).$$

The other equality in (ii) is similar, and (iii) is immediate:

$$e_{j-1}(I)e_{j+1}(I) := e_{j-1}(IR_{\mathfrak{m}})e_{j+1}(IR_{\mathfrak{m}}) \leq e_j(IR_{\mathfrak{m}})^2 =: e_j(I)^2.$$

- (b) For Proposition 2.35 (i), we note that a homogeneous ideal of height n is automatically primary to \mathfrak{m} and appeal to part (a). For (iii), we note that when h_1, \dots, h_{n-j} are homogeneous and sufficiently general, $\text{ht}(I + (h_1, \dots, h_{n-j}))$ is homogeneous of height n and thus \mathfrak{m} -primary. Additionally, we note that $I + \mathfrak{m}^t$ is \mathfrak{m} -primary for all $t > 0$, after which we appeal to part (a) (ii) of this proposition. Similarly to part (a), Proposition 2.37 (iv) follows from the local case. \square

Lemma 2.43. *Suppose we are in one of the following situations:*

- (1) *L/k is a field extension, $R = k[x_1, \dots, x_n]$, $\mathfrak{m} = (x_1, \dots, x_n)$, $S = L[x_1, \dots, x_n]$, and $I \subseteq R$ is \mathfrak{m} -primary or homogeneous.*
- (2) *$(R, \mathfrak{m}) \rightarrow (S, \mathfrak{n})$ is an extension of n -dimensional regular local rings, $\mathfrak{m}S = \mathfrak{n}$, and $I \subseteq R$ is an ideal.*

Then for all $1 \leq j \leq n$ such that $\sigma_j(I)$ is defined, we have $\sigma_j(I) = \sigma_j(IS)$.

Proof. The assumptions imply that $\lambda_R(J) = \lambda_S(JS)$ for every \mathfrak{m} -primary ideal $J \subseteq R$. In particular, this applies to the ideals $(I + \mathfrak{m}^t)^a \mathfrak{m}^b$ for all $a, b, t > 0$; the result follows from Definitions 2.32 and 2.36. \square

3. PROOF OF THEOREM A

3.1. F -Pure Thresholds and the Demailly-Pham Invariant. In this subsection, we require an asymptotic version of [BA17, Theorem 13] in arbitrary characteristic. Our result follows from [BA17; DP14] with the necessary changes being made.

Definition 3.1. Let k be a field and $R = k[x_1, \dots, x_n]$. The **reverse lexicographic order** on R , denoted by $>_{\text{rlex}}$, is the monomial order such that $\mathbf{x}^a > \mathbf{x}^b$ if and only if there exists an index $0 \leq i \leq n-1$ such that $(a_n, \dots, a_{n-i+1}) = (b_n, \dots, b_{n-i+1})$ and $a_{n-i} < b_{n-i}$. The **graded reverse lexicographic order**, denoted $>_{\text{grlex}}$, is the monomial order given by $\mathbf{x}^a >_{\text{grlex}} \mathbf{x}^b$ if and only if $\deg(\mathbf{x}^a) > \deg(\mathbf{x}^b)$ or else $\deg(\mathbf{x}^a) = \deg(\mathbf{x}^b)$ and $\mathbf{x}^a >_{\text{rlex}} \mathbf{x}^b$.

Lemma 3.2. *Let k be a field, $R = k[x_1, \dots, x_n]$, and $J \subseteq R$ an ideal. For $1 \leq j \leq n$, we define $\pi_j : R \rightarrow R/(x_{j+1}, \dots, x_n) \cong k[x_1, \dots, x_j]$. Then*

$$\text{in}_{>_{\text{rlex}}} \pi_j(J) = \pi_j(\text{in}_{>_{\text{rlex}}}(J)).$$

Proof. Let $f \in J$. Write $f = g + h$, where $h \in (x_{j+1}, \dots, x_n)$ and $g \in k[x_1, \dots, x_j]$. If $g = 0$, then $\pi_j(f) = 0$. If $g \neq 0$, then $\text{in}_{>}(f) = \text{in}_{>}(g)$. In both cases, we have $\pi_j(\text{in}_{>}(f)) = \text{in}_{>}(\pi_j(f))$. \square

The following is a general lemma which will be used repeatedly throughout the rest of this article.

Lemma 3.3. *Let L be a field, $S = L[x_1, \dots, x_n]$, and $J \subseteq S$ an \mathfrak{m} -primary homogeneous ideal generated by forms of degree $\leq d$. Then $\mathfrak{m}^d \subseteq \overline{J}$.*

Proof. We first prove the result in the case that L is infinite. First, choose forms f_1, \dots, f_n from among the generators of J such that (f_1, \dots, f_n) is \mathfrak{m} -primary. If h_1, \dots, h_n are general linear forms, then

$$J' := (h_1^{d-\deg(f_1)} f_1, \dots, h_n^{d-\deg(f_n)} f_n)$$

is an \mathfrak{m} -primary (d, \dots, d) -complete intersection contained in J . As $J' \subseteq \mathfrak{m}^d$ and $e(J)' = d^n = e(\mathfrak{m}^d)$, we have $\mathfrak{m}^d = \overline{\mathfrak{m}^d} \subseteq \overline{J'} \subseteq \overline{J}$ by Theorem 2.27.

Now, let L be an arbitrary field, and set $S' = \overline{L}[x_1, \dots, x_n]$. We then have $\overline{J} = \overline{J S'} \cap S \supseteq \mathfrak{m}^d S' \cap S = \mathfrak{m}^d$ by Proposition 2.26 (vii) and the infinite field case. \square

Lemma 3.4 ([BA17], Theorem 4). *Let k be an infinite field, $R = k[x_1, \dots, x_n]$, and I an \mathfrak{m} -primary ideal. If $>$ denotes the monomial order $>_{\text{rlex}}$ and $\gamma \in \text{GL}_n(k)$ is general, then for all $1 \leq j \leq n$ we have*

$$(8) \quad \lim_{t \rightarrow \infty} \frac{e_j(\text{in}_{>}(\gamma^{-1} I^t))}{t^j} = e_j(I).$$

To reassure the reader that Bivià-Ausina's arguments hold in arbitrary characteristic, we summarize the main argument below.

Proof of Lemma 3.4. By Proposition 2.35 (ii), choose $\gamma \in \mathrm{GL}_n(k)$ so that $e_j(I) = e_j\left(\frac{I + \gamma(x_n, \dots, x_{n-j+1})}{\gamma(x_n, \dots, x_{n-j+1})}\right)$. Set $J = \gamma^{-1}I$. By Proposition 2.35 (ii), let $\gamma_t \in \mathrm{GL}_n(k)$ such that $e_j(\mathrm{in}_>(J^t)) = e(\pi_j(\gamma_t^{-1} \mathrm{in}_>(J^t)))$. By [Smi20, Theorem 3.4 (ii)], Hilbert-Samuel multiplicity is upper semicontinuous. Applying this semicontinuity to the specializations $\pi_j(\gamma_t^{-1} \mathrm{in}_>(J^t)) \rightsquigarrow \pi_j(\mathrm{in}_>(J^t))$ and $J^t \rightsquigarrow \mathrm{in}_>(J^t)$ as in [BA17, Proposition 2 and Theorem 4], we deduce

$$(9) \quad e(\pi_j(\mathrm{in}_>(J^t))) \geq e(\pi_j(\gamma_t^{-1} \mathrm{in}_>(J^t))) = e_j(\mathrm{in}_>(J^t)) \geq e_j(J).$$

By [Mus02, Corollary 1.13], Equation (9), and Lemma 3.2 we have

$$(10) \quad \begin{aligned} e_j(I) &= e_j(J) = e(\pi_j(J)) = \lim_{t \rightarrow \infty} \frac{e(\pi_j(J)^t)}{t^j} \\ &= \lim_{t \rightarrow \infty} \frac{e(\mathrm{in}_>(\pi_j(J^t)))}{t^j} = \lim_{t \rightarrow \infty} \frac{e(\pi_j(\mathrm{in}_>(J^t)))}{t^j} \\ &\geq \lim_{t \rightarrow \infty} \frac{e_j(\mathrm{in}_>(J^t))}{t^j} \geq \lim_{t \rightarrow \infty} \frac{e_j(J^t)}{t^j} = e_j(J). \end{aligned}$$

It follows that the inequalities in Equation (10) must be sharp, proving the claim. \square

Remark 3.5. The generic change of coordinates in Lemma 3.4 is necessary when $\dim(R) \geq 3$. Consider $R = \mathbb{Q}[x, y, z]$ and $I = (xy, x^2 + z^2, x^4, y^4, z^4)$. Let \succ denote the graded reverse lexicographic order with $x \succ y \succ z$. Using the ReesAlgebra package in Macaulay2 [Eis+; GS], we compute $e(\mathrm{in}_\succ(I^5)) = 2000$. Separately, we have $2000 = 5^3 \cdot e(I) = e(I^5)$. For all $t > 0$, we have $\mathrm{in}_\succ(I^{5t}) \subseteq \mathrm{in}_\succ(I^5)^t$, hence

$$t^3 e(I^5) = e(I^{5t}) \leq e(\mathrm{in}_\succ(I^{5t})) \leq e(\mathrm{in}_\succ(I^5)^t) = t^3 e(I^5),$$

Set $\mathfrak{a} = \mathrm{in}_\succ(I^5)$. By Theorem 2.27, we deduce that $\overline{\mathrm{in}_\succ(I^{5t})} = \overline{\mathfrak{a}^t}$ for all $t > 0$. It follows from Proposition 2.39 that

$$\lim_{t \rightarrow \infty} \frac{e_2(\mathrm{in}_\succ(I^t))}{t^2} = \lim_{t \rightarrow \infty} \frac{e_2(\mathrm{in}_\succ(I^{5t}))}{(5t)^2} = \frac{e_2(\mathfrak{a}^t)}{(5t)^2} = \frac{e_2(\mathfrak{a})}{5^2}.$$

To compute $e_2(\mathfrak{a})$, we use Proposition 2.35 (ii). Suppose $h = ax + by + cz$ is such that $e_2(\mathfrak{a}) = e\left(\frac{\mathfrak{a} + (h)}{(h)}\right)$. Let $\gamma \in \mathrm{GL}_n(k)$ denote the map $(x, y, z) \mapsto (ax, by, cz)$. As \mathfrak{a} is a monomial ideal, we have $\gamma^{-1}\mathfrak{a} = \mathfrak{a}$, hence

$$(11) \quad e\left(\frac{\mathfrak{a} + \gamma(x + y + z)}{\gamma(x + y + z)}\right) = e\left(\frac{\mathfrak{a} + (x + y + z)}{(x + y + z)}\right).$$

The quantity in Equation (11) can be computed directly using the ReesAlgebra package, giving us

$$\lim_{t \rightarrow \infty} \frac{e_2(\mathrm{in}_\succ(I^t))}{t^2} = \frac{e_2(\mathrm{in}_\succ(I^5))}{5^2} = \frac{22}{5}.$$

Using Lemmas 3.7 and 3.8, we compute $e_2(I) = 4 < \frac{22}{5}$.

Definition 3.6. Let k be a field, $R = k[x_1, \dots, x_n]$, $\mathfrak{m} = (x_1, \dots, x_n)$, and let \mathbf{a}_\bullet be a graded system of \mathfrak{m} -primary ideals. We define:

- The asymptotic mixed multiplicities: $e_j(\mathbf{a}_\bullet) = \liminf_m \frac{e_j(\mathbf{a}_m)}{m^j}$.
- The asymptotic Demailly-Pham invariant: $E_n(\mathbf{a}_\bullet) = \frac{1}{e_1(\mathbf{a}_\bullet)} + \dots + \frac{e_{n-1}(\mathbf{a}_\bullet)}{e_n(\mathbf{a}_\bullet)}$
- The asymptotic singularity threshold: $c(\mathbf{a}_\bullet) = \liminf_m mc(\mathbf{a}_m)$.

Before we prove Theorem 3.9 and our asymptotic version of [BA17, Theorem 13], we require the following standard facts.

Lemma 3.7. *Let L be a field and $S = L[x_1, \dots, x_n]$. Let I be a homogeneous ideal of S and $J \subseteq I$ denote the ideal of S generated by the homogeneous forms in I of degree $\leq d$. If $\text{ht}(J) = n$, then $\bar{J} = \bar{I}$.*

Proof. It is clear that $\bar{J} \subseteq \bar{I}$. Let $\mathfrak{m} := (x_1, \dots, x_n)$. For the reverse containment, note that $I \subseteq J + \mathfrak{m}^{d+1}$. By Lemma 3.3 we have

$$\bar{I} \subseteq \overline{J + \mathfrak{m}^{d+1}} \subseteq \overline{J + \mathfrak{m}^d} \subseteq \bar{J}.$$

□

Lemma 3.8. *Let L be a field and $S = L[x_1, \dots, x_n]$. Let $J = (f_1, \dots, f_n)$ be a complete intersection where $\deg f_i = d_i$ and $d_1 \leq \dots \leq d_n$. Then we have the following:*

- (i) *If L is infinite, then for a general hyperplane section $H \subseteq \text{Spec } S$, we have $e(J|_H) = d_1 \cdots d_{n-1}$.*
- (ii) *Let $1 \leq j \leq n$. With no assumption on $|L|$, we have $e_j(J) = d_1 \dots d_j$, and hence $E_j(J) = \frac{1}{d_1} + \dots + \frac{1}{d_j}$.*

Proof. As the f_1, \dots, f_n form a homogeneous system of parameters for S , we have

$$(12) \quad e(J) = d_1 \dots d_n.$$



Choosing H generally so that $\text{ht}(f_1, \dots, f_{n-1})|_H = n-1$, Lemma 3.7 gives $e(J|_H) = e((f_1|_H, \dots, f_{n-1}|_H))$, so (i) follows from Equation (12). For (ii), Lemma 2.43 shows that $e_j(J)$ is invariant under extension of the base field, so it suffices to consider the case of an infinite field. The result follows from (i) and Proposition 2.35 (ii). □

Theorem 3.9. *Let n be a positive integer and $1 \leq l \leq n$. Suppose either:*

- (1) *(R, \mathfrak{m}) is a regular local ring of dimension n , excellent if $\text{char } R = 0$, and that $I \subseteq R$ is an ideal of height at least l ;*
- (2) *k is a field, $R = k[x_1, \dots, x_n]$, $\mathfrak{m} = (x_1, \dots, x_n)$, and $I \subseteq R$ is an ideal which is \mathfrak{m} -primary or homogeneous of height at least l .*

Then $E_l(I) \leq c(I)$.

Proof. We reduce to the case of an \mathfrak{m} -primary ideal in a polynomial ring over an infinite field. In the case (2), Proposition 2.10 (vi) and Lemma 2.43 show that E_l, c are invariant under field extension. If J is an ideal generated by $n-l$ general linear forms, then $E_l(\frac{I+J}{J}) = E_l(I)$ and $c(\frac{I+J}{J}) \leq c(I)$ by Proposition 2.10 (iv) and Proposition 2.42, so we may assume $l = n$.


 FIGURE 2. $\Gamma(\mathfrak{a}_\bullet)$, together with $\mu\mathbf{1}$ and H .

 FIGURE 3. $\Gamma(\mathfrak{a}'_\bullet)$ in terms of H .

Suppose we are in the case (1). By Proposition 2.10 (vi) and Lemma 2.43, the quantities c, E_l are invariant under completion and field extension, so we may without loss of generality assume R is complete local with infinite residue field k . As $c(I) \geq c((I+J)/J)$ for any ideal $J \subseteq R$, we let J be an ideal generated by $n-l$ elements of \mathfrak{m} whose images in $\mathfrak{m}/\mathfrak{m}^2$ are sufficiently general; it suffices to show that $c((I+J)/J) \geq E_l((I+J)/J)$, where $(I+J)/J$ is an ideal of height l in a complete regular local ring of dimension l . Therefore, without loss of generality, we may assume $l = n$, and hence I is \mathfrak{m} -primary. Set $R' = k[x_1, \dots, x_n], \mathfrak{m}' = (x_1, \dots, x_n)$. Since I is \mathfrak{m} -primary, there exists $I' \subseteq R'$ with $I'R = I$. Another application of Proposition 2.10 (vi) and Lemma 2.43 gives $c(I) = c(I'R'_{\mathfrak{m}'})$ and $E_n(I) = E_n(I'R'_{\mathfrak{m}'})$, so by Notation 2.9 and Definition 2.40 we have $c(I) = c(I'), E_n(I) = E_n(I')$.

Let $\gamma \in \mathrm{GL}_n(k)$ such that Equation (8) holds. Set $\mathfrak{a}_m = \mathrm{in}_{>}(\gamma^{-1}I^m)$ for $m > 0$. By Lemma 3.4 we have $E_n(\mathfrak{a}_\bullet) = E_n(I)$ and by Proposition 2.22 we have $c(\mathfrak{a}_\bullet) \leq c(I)$. Let $\mu = \inf\{t : t\mathbf{1} \in \Gamma\}$. Since Γ is convex and $\mu\mathbf{1} \in \partial\Gamma$, by [Roc70, Corollary 11.6.1] there exists a half-space $H^+ \subseteq \mathbb{R}^n$ such that $\Gamma \subseteq H^+$ and that $\mu\mathbf{1} \in \partial H^+$. Since Γ is closed under translation by elements of $\mathbb{R}_{\geq 0}^n$ and the complement of Γ in $\mathbb{R}_{\geq 0}^n$ is bounded, the same is true for H^+ . The closure of the complement of H^+ in $\mathbb{R}_{\geq 0}^n$ is therefore a simplex, which we denote by $\mathrm{conv}(\mathbf{0}, (a_1, 0, \dots, 0), \dots, (0, \dots, 0, a_n))$. See Figures 2 and 3 for a depiction of this step.

Define a graded system of monomial ideals \mathfrak{a}_\bullet by $\mathfrak{a}'_m = \{\mathbf{x}^{\mathbf{a}} : \mathbf{a} \in mH^+\}$. By assumption that $\Gamma \subseteq H^+$, we have $\mathfrak{a}_m \subseteq \mathfrak{a}'_m$ for all m . Consequently, we have $E_n(\mathfrak{a}_\bullet) \leq E_n(\mathfrak{a}'_\bullet)$ by Proposition 2.39. By Proposition 2.16, we also have $c(\mathfrak{a}_\bullet) = c(\mathfrak{a}'_\bullet)$. We can compute $E_n(\mathfrak{a}'_\bullet)$ from the numbers a_1, \dots, a_n : by observing that

$$\overline{(x_1^{\lfloor ma_1 \rfloor}, \dots, x_n^{\lfloor ma_n \rfloor})} \subseteq \mathfrak{a}'_m \subseteq \overline{(x_1^{\lfloor ma_1 \rfloor}, \dots, x_n^{\lfloor ma_n \rfloor})},$$

it follows from Lemma 3.8 that $E_n(\mathfrak{a}'_\bullet) = \frac{1}{a_1} + \cdots + \frac{1}{a_n}$. All together, we have

$$E_n(I) = E_n(\mathfrak{a}_\bullet) \leq E_n(\mathfrak{a}'_\bullet) = \frac{1}{a_1} + \cdots + \frac{1}{a_n} = c(\mathfrak{a}'_\bullet) = c(\mathfrak{a}_\bullet) \leq c(I).$$

□

Remark 3.10. The insight depicted in Figures 2 and 3 appears earlier than the work of Demailly, Pham and Bivià-Ausina, in the proof [FEM04, Theorem 1.1].

Corollary 3.11. *Let k be an uncountably infinite field, $R = k[x_1, \dots, x_n]$, and I an \mathfrak{m} -primary homogeneous ideal. Let $>$ denote the graded reverse lexicographic order and let $\mathfrak{a}_m := \text{gin}(I^m)$. Lastly, let $\mathbf{b}_1, \dots, \mathbf{b}_n$ denote the standard unit vectors in \mathbb{R}^n . If $E_n(I) = c(I)$, then*

$$(13) \quad \overline{\mathbb{R}_{\geq 0}^n \setminus \Gamma(\mathfrak{a}_\bullet)} = \text{conv} \left(\mathbf{0}, e_1(I)\mathbf{b}_1, \frac{e_2(I)}{e_1(I)}\mathbf{b}_2, \dots, \frac{e_n(I)}{e_{n-1}(I)}\mathbf{b}_n \right).$$

Proof. By construction, $>$ agrees with the reverse lexicographic order on homogeneous ideals. Consequently, we may choose $\gamma \in \text{GL}_n(k)$ very generally so that Equation (8) holds and $\text{in}_>(\gamma^{-1}I^m) = \text{gin}_>(I^m)$ for all $m > 0$. With this choice of γ , let $\mathfrak{a}_\bullet, \mathfrak{a}'_\bullet$ be as in Theorem 3.9.

Suppose $E_n(I) = c(I)$. Then we also have $E_n(\mathfrak{a}_\bullet) = E_n(\mathfrak{a}'_\bullet)$. By [BA17, Proposition 10], we further have that $e_j(I) = e_j(\mathfrak{a}_\bullet) = e_j(\mathfrak{a}'_\bullet)$ for all $1 \leq j \leq n$. In particular, $e_n(\mathfrak{a}_\bullet) = e_n(\mathfrak{a}'_\bullet)$, so by [Mus02, Theorem 2.12 and Lemma 2.13], we have $\text{vol}(\mathbb{R}_{\geq 0}^n \setminus \Gamma(\mathfrak{a}_\bullet)) = \text{vol}(\mathbb{R}_{\geq 0}^n \setminus \Gamma(\mathfrak{a}'_\bullet))$. Since $\Gamma(\mathfrak{a}_\bullet), \Gamma(\mathfrak{a}'_\bullet)$ are closed and convex with positive volume, it follows that $\Gamma(\mathfrak{a}_\bullet) = \Gamma(\mathfrak{a}'_\bullet)$. Since the generic initial ideal is Borel-fixed, we have $a_1 \leq \cdots \leq a_n$, hence $e_j(\mathfrak{a}'_\bullet) = a_1 \dots a_j$ for all $1 \leq j \leq n$, proving the claim. □

Remark 3.12. Equation (13) is a necessary condition to have $c(I) = E_n(I)$, but not a sufficient condition. By [May14] in characteristic zero, Equation (13) holds for any homogeneous complete intersection $J = (f_1, \dots, f_n)$.

In fact, Equation (13) holds for any homogeneous complete intersection in positive characteristic, though a small modification to Mayes's argument in [May14] must be made. In particular, Lemma 3.6 in op. cit. does not hold in positive characteristic: in the ring $\overline{\mathbb{F}}_p[x, y]$ we have $\text{gin}(x^p, y^p) = (x^p, y^p)$. Instead, one appeals to Theorem 2.29. The details will appear in the author's thesis.

Example 3.13 ([Fer26]). Here we give a counterexample to [Kim21, Question 1.3], which also functions as a counterexample to the analogous statement in positive characteristic. Let k be a field and set $R = k[x, y, z]$. The idea is to start with an ideal $\mathfrak{c} \subseteq R$ and a general hyperplane H for which the equality $c(\mathfrak{c}) - c(\mathfrak{c}|_H) = \frac{e_2(\mathfrak{c})}{e_3(\mathfrak{c})}$ holds and to produce an ideal $\mathfrak{a} \supseteq \mathfrak{c}$ for which $c(\mathfrak{a}) = c(\mathfrak{c}), c(\mathfrak{a}|_H) = c(\mathfrak{c}|_H), e_2(\mathfrak{a}) = e_2(\mathfrak{c})$, and $e(\mathfrak{a}) < e(\mathfrak{c})$.

We'll define a family of monomial ideals satisfying the desired conditions, which we depict in Figures 4 and 5. Let $3 \leq a < b$. We define the auxiliary ideal $\mathfrak{b} =$




FIGURE 4. $\Gamma(\mathfrak{c})$, with $\mu\mathbf{1}$ computing $c(\mathfrak{c})$ in red.

FIGURE 5. $\Gamma(\mathfrak{a})$, with $\mu\mathbf{1}$ computing $c(\mathfrak{a})$ in red.

(x^a, xy, y^a) . Set $\mathfrak{c} = (\mathfrak{b}, z^b)$ and $\mathfrak{a} = (\mathfrak{c}, zx^{a-1})$. As $a \geq 3$, we have $\mathfrak{a} \subseteq \overline{(x^2, y^2, z^b)}$. Using Proposition 2.16, we compute

$$1 + \frac{1}{b} = c((xy, z^b)) \leq c(\mathfrak{c}) \leq c(\mathfrak{a}) \leq c(\overline{(x^2, y^2, z^b)}) = 1 + \frac{1}{b}.$$

As in Remark 3.5, the linear form $h = x - y - z$ is sufficiently general to compute $e_2(\mathfrak{a})$ and $\max_\ell c(\mathfrak{a}|_{\ell=0})$. Identifying R/hR with $k[x, y]$ and letting $\mathfrak{a}', \mathfrak{b}', \mathfrak{c}'$ denote the images of $\mathfrak{a}, \mathfrak{b}, \mathfrak{c}$ in R/hR , we have

$$\mathfrak{b}' \subseteq \mathfrak{c}' \subseteq \mathfrak{a}' = (\mathfrak{b}', (x+y)x^{a-1}, (x+y)^b) \subseteq \overline{\mathfrak{b}'},$$

so we have $\overline{\mathfrak{a}'} = \overline{\mathfrak{b}'} = \overline{\mathfrak{c}'}$, which implies that $e_2(\mathfrak{a}) = e_2(\mathfrak{c})$ and $c(\mathfrak{a}') = c(\mathfrak{c}')$.

By Lemma 3.3 and Lemma 3.8 respectively, we have $\mathfrak{b}' = \overline{(xy, x^a + y^a)}$ and $e(\mathfrak{b}') = 2a$. A similar computation yields $\mathfrak{c} = \overline{(xy, x^a + y^a, z^b)}$, so $e(\mathfrak{c}) = 2ab$. On the other hand, we claim that the containment $\overline{\mathfrak{c}} \subseteq \overline{\mathfrak{a}}$ is strict — in particular, we claim that $zx^{a-1} \notin \overline{\mathfrak{c}}$. To see this, we note that the monomials in $k[x, z] \cap \overline{\mathfrak{c}}$ are precisely those $x^i z^j$ for which $i/a + j/b \geq 1$, and we have $(a-1)/a + 1/b < 1$. By Theorem 2.27, it follows that $e(\mathfrak{a}) < e(\mathfrak{c})$. We deduce that

$$c(\mathfrak{a}) = 1 + \frac{1}{b} = c(\mathfrak{c}) + \frac{e_2(\mathfrak{c})}{e(\mathfrak{c})} < c(\mathfrak{a}|_{h=0}) + \frac{e_2(\mathfrak{a})}{e(\mathfrak{a})},$$

violating the conjectured inequality.

In light of [Kim21, Proposition 2.4 (i)], we might instead consider the weaker bound $c(I) - c(I|_H) \geq \frac{1}{\mathcal{L}_0(I)}$, where $\mathcal{L}_0(I)$ is the Łojasiewicz exponent (Definition 4.5). This does hold; see Lemma 4.6.

4. THEOREM B IN THE COMPLETE INTERSECTION CASE

4.1. Behavior of the Singularity Threshold Under Modification. Fix the following notation throughout this subsection.

Assumption 4.1. *Let k be a field, $R = k[x_1, \dots, x_n]$, and let \mathfrak{m} denote the homogeneous maximal ideal. Let $I \subseteq R$ be an \mathfrak{m} -primary homogeneous ideal. Write $I = I_1 + \dots + I_r$, where I_j is generated by forms of degree d_j and $d_1 < \dots < d_r$.*

Lemma 4.2 ([BS15], Lemma 3.2). *Let $R = k[x_1, \dots, x_n]$ and let \mathfrak{m} denote the homogeneous maximal ideal. For any $e, t \in \mathbb{Z}^+$, we have*

$$(\mathfrak{m}^{[p^e]} : \mathfrak{m}^t) = \begin{cases} R & t \geq np^e - n + 1 \\ \mathfrak{m}^{[p^e]} + \mathfrak{m}^{np^e - n + 1 - t} & t < np^e - n + 1 \end{cases}$$

More generally, we have the following.

Lemma 4.3. *Let $R = k[x_1, \dots, x_n]$. Let v be a monomial valuation on R with $v(x_i) \geq 0$ for all $1 \leq i \leq n$. For $\lambda \in \mathbb{R}^+$, let \mathfrak{a}_λ denote the ideal $\{f \in R : v(f) \geq \lambda\}$ and $\mathfrak{a}_\lambda^+ = \{f \in R : v(f) > \lambda\}$. Let $q \in \mathbb{Z}^+$, $\lambda \in \mathbb{R}^+$. Then we have*

$$(14) \quad ((x_1^q, \dots, x_n^q) : \mathfrak{a}_\lambda) = (x_1^q, \dots, x_n^q) + \mathfrak{a}_{(q-1)v(x_1 \dots x_n) - \lambda}^+$$

Proof. The argument is the same as Lemma 4.2. Let $m \notin (x_1^q, \dots, x_n^q)$ be a monomial. Then $m \mid (x_1 \dots x_n)^{q-1}$, so $\mathfrak{a}_\lambda m \not\subseteq (x_1^q, \dots, x_n^q)$ if and only if $\frac{(x_1 \dots x_n)^{q-1}}{m} \in \mathfrak{a}_\lambda$, which holds if and only if $v((x_1 \dots x_n)^{q-1}) - v(m) \leq \lambda$. We've shown that the two sides of Equation (14) contain the same monomials; both sides are monomial ideals, so the result follows. \square

Lemma 4.4 ([Bai25], Lemma 4.2). *Let k be a field of characteristic $p > 0$, let $R = k[x_1, \dots, x_n]$, and $I \subseteq R$ a homogeneous ideal. For a hyperplane H cut out by a linear form ℓ , we let $I|_H$ denote the image of I in $R/\ell R$. In this case, we have*

$$\nu_{I|_H}(p^e) \leq \max\{r : I^r \not\subseteq \mathfrak{m}^{[p^e]} + \mathfrak{m}^{(n-1)(p^e-1)+1}\},$$

Definition 4.5. Let $R = k[x_1, \dots, x_n]$ and let $I \subseteq R$ be an ideal. The **Łojasiewicz exponent** of I , denoted $\mathcal{L}_0(I)$, is equal to the least integer t such that $\mathfrak{m}^t \subseteq \bar{I}$, or else ∞ in the case that I is not \mathfrak{m} -primary.

Lemma 4.6. *Let $R = k[x_1, \dots, x_n]$ and let $I \subseteq R$ be an ideal. If $H \subseteq \text{Spec } R$ is a hyperplane, then we have $c(I) - c(I|_H) \geq \frac{1}{\mathcal{L}_0(I)}$.*

Proof. If $\mathcal{L}_0(I) = \infty$ then there is nothing to prove, so suppose $\mathcal{L}_0(I) = d < \infty$.

We first prove the claim in characteristic $p > 0$. Combining Lemma 4.2 and Lemma 4.4, we have

$$\nu_{I|_H}(p^e) \leq \max\{s : \mathfrak{m}^{p^e} I^s \not\subseteq \mathfrak{m}^{[p^e]}\}.$$

Consequently, we have $\max\{s : \mathfrak{m}^{p^e} I^s \not\subseteq \mathfrak{m}^{[p^e]}\} \leq \nu_I(p^e) - \left\lfloor \frac{p^e}{d} \right\rfloor$, which implies that $\nu_{I|_H}(p^e) \leq \nu_I(p^e) - \left\lfloor \frac{p^e}{d} \right\rfloor$. Dividing by p^e and taking the limit as $e \rightarrow \infty$ gives the result.

We now prove the claim in characteristic 0. The result follows from Proposition 2.8, the details of which we spell out explicitly in this case (though moving forward, we will be more terse). Let $A \subseteq k$ be a finitely-generated \mathbb{Z} -algebra and $J \subseteq A[x_1, \dots, x_n]$ an ideal such that $JR = I$. Such a subring A can always be constructed by adjoining to \mathbb{Z} the field coefficients appearing in a generating set for I ; we'll also choose A to contain the coefficients of the linear form ℓ . If μ is a maximal ideal of A , we let I_μ denote the image of J in $(A/\mu)[x_1, \dots, x_n]$. Applying Proposition 2.8 to both I and $I|_H$, we obtain

$$c(I) - c(I|_H) = \lim_{\substack{\mu \in \text{Spec } A \\ \text{char } A/\mu \rightarrow \infty}} c(I|_\mu) - c(I_\mu|_{H_\mu}) \geq 1/d.$$

□

Corollary 4.7. *Assume the setup of Assumption 4.1 and let $H \subseteq \text{Spec } R$ be a hyperplane cut out by a linear form ℓ . Then $c(I) - c(I|_H) \geq 1/d_r$.*

Proof. By Lemma 3.3, we have $\mathfrak{m}^{d_r} \subseteq \bar{I}$, so $\mathcal{L}_0(I) \leq d_r$. The result follows from Lemma 4.6. □

Lemma 4.8. *Assume the setup of Assumption 4.1. Suppose $r = 2$. Then we have $c(I) = \frac{n}{d_2} + c(I_1) \frac{d_2 - d_1}{d_2}$. In particular, for any $0 \leq s \leq n$, we have $c(I) = \frac{s}{d_1} + \frac{n-s}{d_2}$ if and only if $c(I_1) = \frac{s}{d_1}$.*

Proof. We prove the claim first in positive characteristic. By Lemma 3.3, we have $\mathfrak{m}^{d_2} \subseteq \bar{I}$, so $I \subseteq I_1 + \mathfrak{m}^{d_2} \subseteq \bar{I}$, so $c(I) = c(I_1 + \mathfrak{m}^{d_2})$. Consequently, we have

$$\nu_{\bar{I}}(p^e) = \max \left\{ a + b : I_1^a \mathfrak{m}_\mu^{bd_n} \not\subseteq \mathfrak{m}^{[p^e]} \right\} = \max \{ a + b : I_1^a \not\subseteq (\mathfrak{m}_p^{[p^e]} : \mathfrak{m}_p^{bd_n}) \}.$$

By Lemma 4.2, we obtain

$$(15) \quad \nu_{\bar{I}}(p^e) = \max \{ a + b : I_1^a \not\subseteq \mathfrak{m}^{[p^e]} + \mathfrak{m}^{np^e - n + 1 - bd_2} \}.$$

The ideal I_1^a is generated in degree ad_1 , so $I_1^a \not\subseteq \mathfrak{m}^{[p^e]} + \mathfrak{m}^t$ if and only if $I_1^a \not\subseteq \mathfrak{m}^{[p^e]}$ and $ad_1 < t$. With this insight, we refine Equation (15):

$$(16) \quad \nu_I(p^e) = \max_{0 \leq a \leq \nu_{I_1}(p^e)} a + \frac{np^e - n - ad_1}{d_2}.$$

The quantity being maximized in Equation (16) is an increasing function of a , so the maximum occurs at $a = \nu_{I_1}(p^e)$ and

$$\nu_{\bar{I}}(p^e) = \frac{np^e - n}{d_2} + \nu_{I_1}(p^e) \frac{d_2 - d_1}{d_2}.$$

Dividing by p^e and letting $e \rightarrow \infty$, we obtain

$$c(I) = \frac{n}{d_2} + c(I_1) \frac{d_2 - d_1}{d_2}.$$

For characteristic zero, we can spread out to positive characteristic as in Corollary 4.7 and compute

$$c(I) = \lim_{\mu} c(I_{\mu}) = \lim_{\mu} \left(\frac{n}{d_2} + c(I_{1,\mu}) \frac{d_2 - d_1}{d_2} \right) = \frac{n}{d_2} + c(I_1) \frac{d_2 - d_1}{d_2}.$$

□

4.2. Induction Setup. In this subsection, we begin the proof of Theorem 5.1 in the case that I is a complete intersection and $k = \bar{k}$.

Assumption 4.9. We assume a setup similar to Assumption 4.1. Let k be an algebraically closed field. Let $a_1, \dots, a_r, d_1, \dots, d_r \in \mathbb{Z}^+$. For $1 \leq i \leq r$, let \mathbf{x}_i denote the tuple of variables $x_{i,1}, \dots, x_{i,a_i}$, and let $R = k[\mathbf{x}_1, \dots, \mathbf{x}_r]$. Let $n = a_1 + \dots + a_r = \dim R$. Let $I \subseteq R$ be a complete intersection of the form $(f_{1,1}, \dots, f_{r,a_r})$ such that $f_{i,j}$ is a d_i -form. For $1 \leq j \leq r$, write $I_j = (f_{j,1}, \dots, f_{j,a_j})$. Let w_d denote the monomial valuation with $w_d(x_{i,j}) = 1/d_i$. We let \mathfrak{D} denote the ideal $\overline{(\mathbf{x}_1^{d_1}, \dots, \mathbf{x}_r^{d_r})}$, which coincides with the set of elements z with $w_d(z) \geq 1$.

Assumption 4.10. Assume the setup of Assumption 4.9. We define the following condition on the ideal I :

$$(\dagger) \quad I_1 \text{ is extended from } k[\mathbf{x}_1] \text{ and } I \subseteq \mathfrak{D} + (\mathbf{x}_1)$$

Definition 4.11. For $r \in \mathbb{Z}^+$, we define the statements A_r, B_r .

$$(A_r) \quad \text{For all } I \text{ as in Assumption 4.9, if } E_n(I) = c(I) \text{ then} \\ \text{there exists } \gamma \in \text{GL}_n(k) \text{ such that } \gamma I \subseteq \mathfrak{D}.$$

$$(B_r) \quad \text{For all } I \text{ as in Assumption 4.9, if } E_n(I) = c(I) \text{ then} \\ \text{there exists } \gamma \in \text{GL}_n(k) \text{ such that } \gamma I \text{ satisfies Equation } (\dagger).$$

The goal of this section is to prove A_r for all r . We accomplish this via the following steps:

- (1) A_1, A_2 hold;
- (2) For $r \geq 3$, A_{r-1} combined with A_2 implies B_r ;
- (3) For $r \geq 3$, B_r implies A_r .

The proof of step (1) is short, but steps (2) and (3) will each have a dedicated subsection.

Proof of A_1 . If I is an \mathfrak{m} -primary complete intersection generated by forms of degree d for some d , by Lemma 3.3 we have $\bar{I} = \mathfrak{m}^d$. □

Remark 4.12. When $r = 1$, the above argument shows that $c(I) = E_n(I)$ is satisfied for all choices of I . Unfortunately, this means that the proof of our induction step $A_2, A_{r-1} \implies A_r$ can't be modified to show $A_1 \implies A_2$. We instead expand the scope of our base case.

To prove A_2 , we introduce the notion of *essential dimension*.

Definition 4.13 (Essential Dimension). Let $J \subseteq R = k[x_1, \dots, x_d]$ be a homogeneous ideal. The essential dimension $\text{ess}(J)$ is equal to the minimal r for which there exist linear forms ℓ_1, \dots, ℓ_r such that J is extended from $I \subseteq k[\ell_1, \dots, \ell_r]$.

We have the following result.

Proposition 4.14 ([Bai25], Proposition 3.3). *Let k be an algebraically closed field, $R = k[x_0, \dots, x_n]$, and $J \subseteq R$ a homogeneous ideal. Set $r = \text{ht}(J)$. Let $L = (\ell_{r+1}, \dots, \ell_n)$, where the ℓ_i are chosen generally. For $r \leq t \leq n$, set $L_t = (\ell_{t+1}, \dots, \ell_n)$ and $J_t = \frac{J+L_t}{L_t}$. Then for all $r \leq t \leq n$, we have $\text{ess}(J_t) = \max(t+1, \text{ess}(J))$.*

Proof of A_2 . By Lemma 4.8, if $c(I) = E_n(I)$ then $c(I_1) = a_1/d_1$. By [FEM03, Theorem 3.5] in characteristic zero and [Bai25, Theorem 3.17] in positive characteristic, it follows that $\text{ess}(I_1) = a_1$. By [Bai25, Lemma 3.18], there exists $\gamma \in \text{GL}_n(k)$ such that $\overline{\gamma(I_1)} = (\mathbf{x}_1)^{d_1}$, hence $\gamma I \subseteq (\mathbf{x}_1)^{d_1} + \mathfrak{m}^{d_2} \subseteq \mathfrak{D}$. \square

4.3. Step (2): For $r \geq 3$, A_{r-1} combined with A_2 implies B_r .

Lemma 4.15. *Assume the setup of Assumption 4.9 and suppose $r \geq 3$, $c(I) = E_n(I)$. If A_{r-1} holds, there exists $\gamma \in \text{GL}_n(k)$ such that $\gamma \overline{I_1} = (\mathbf{x}_1)^{d_1}$.*

Proof. Note that $E_n(I) = a_1/d_1 + \dots + a_r/d_r$ by Lemma 3.8. Let L be an ideal of R generated by $a_3 + \dots + a_d$ general linear forms. Since I is a complete intersection, $I_1 + I_2 + L$ is $\frac{\mathfrak{m}}{L}$ -primary, so by Lemma 3.7 we have $c(R/L, \frac{I+L}{L}) = c(R/L, \frac{I_1+I_2+L}{L})$. Consequently, by repeated application of Corollary 4.7, we have

$$(17) \quad c(I) \geq c\left(R/L, \frac{I_1 + I_2 + L}{L}\right) + \frac{a_3}{d_3} + \dots + \frac{a_r}{d_r}.$$

Assuming $c(I) = E_n(I)$, we have

$$\frac{a_1}{d_1} + \frac{a_2}{d_2} \leq c\left(R/L, \frac{I_1 + I_2 + L}{L}\right) \leq \frac{a_1}{d_1} + \frac{a_2}{d_2},$$

where the left-hand side is by Theorem 3.9 and the right-hand side is by Equation (17). Both inequalities are therefore equalities, so by the proof of A_2 , we have $\text{ess}\left(\frac{I_1+L}{L}\right) = a_1$. By Proposition 4.14, it follows that $\text{ess}(I_1) = a_1$. The result then follows from [Bai25, Lemma 3.18]. \square

Lemma 4.16. *Assume the setup of Assumption 4.9. Suppose $c(I) = E_n(I)$. Then there exists $\gamma \in \text{GL}_n(k)$ such that γI satisfies Equation (\dagger) .*

Proof. By Lemma 4.15, we may assume I_1 is extended from $k[\mathbf{x}_1]$. Let \succ denote the monomial partial order induced by the monomial valuation $w(x_{1,i}) = 0$ and $w(x_{i,j}) = 1$ for $i \geq 2$. For $2 \leq i \leq r, 1 \leq j \leq a_i$, let $g_{i,j} := \text{in}_\succ(f_{i,j})$. Since I is a complete intersection, we have $f_{i,j} \notin \sqrt{I_1} = (\mathbf{x}_1)$, hence $g_{i,j} \notin (\mathbf{x}_1)$ and moreover $f_{i,j} - g_{i,j} \in (\mathbf{x}_1)$. Observe that

$$(18) \quad \text{in}_\succ(I) \supseteq I_1 + \text{in}_\succ(I_2 + \dots + I_r) \supseteq I_1 + (g_{2,1}, \dots, g_{r,a_r}).$$

Let I' denote the right-hand side of Equation (18). Because $g_{i,j}$ and $f_{i,j}$ have the same image modulo $(\mathbf{x}_1) = \sqrt{I_1}$, the ideal I' is a complete intersection. In particular,

I' is a complete intersection of type $(\underbrace{d_1, \dots, d_1}_{a_1}, \dots, \underbrace{d_r, \dots, d_r}_{a_r})$. By Lemma 3.8 and

Proposition 2.22, we have

$$(19) \quad E_n(I) = E_n(I') \leq c(I') \leq c(\text{in}_>(I)) \leq c(I) = E_n(I).$$

As $\bar{I}_1 = (\mathbf{x}_1)^{d_1}$, we have $c(I_1) = a_1/d_1$. Since I_1 and $J := (g_{2,1}, \dots, g_{r,a_r})$ are defined in terms of disjoint sets of variables, we have by [TXZ24, Theorem 2.4 (1)] that

$$(20) \quad c(R, I') = c(k[\mathbf{x}_1], I_1) + c(k[\mathbf{x}_2, \dots, \mathbf{x}_r], J) = \frac{a_1}{d_1} + c(k[\mathbf{x}_2, \dots, \mathbf{x}_r], J).$$

It follows from Equations (19) and (20) that J , which is a complete intersection in $k[\mathbf{x}_2, \dots, \mathbf{x}_r]$, satisfies $E_{a_2+\dots+a_r}(J) = c(J)$. By A_{r-1} , there exists $\gamma \in \text{GL}_{n-a_1}(k)$ such that $\gamma J \subseteq (\mathbf{x}_2^{d_2}, \dots, \mathbf{x}_r^{d_r})$.

We define $\gamma' := \begin{bmatrix} \text{id}_{a_1} & 0 \\ 0 & \gamma \end{bmatrix}$ and we claim that $\gamma'I$ satisfies Equation (†). By construction, $\gamma'I'$ satisfies Equation (†). Since $g_{i,j} - f_{i,j} \in (\mathbf{x}_1)$ for all $2 \leq i \leq r, 1 \leq j \leq a_i$, we have

$$\gamma'(I_2 + \dots + I_r) \subseteq \gamma'(g_{1,1}, \dots, g_{r,a_r}) + (\mathbf{x}_1),$$

which proves that $\gamma'I$ also satisfies Equation (†). \square

Remark 4.17. In the proof of Lemma 4.16, we used the assumption A_{r-1} to show that $I|_H \subseteq \mathfrak{D}|_H$. This condition is necessary, but not sufficient, to have $I \subseteq \mathfrak{D}$ — even if I also satisfies Equation (†). For example, consider the ideal $I = (x^2, y^3 + xyz, z^7)$ and $\mathfrak{D} = (x^2, y^3, z^7)$.

4.4. Step (3): B_r implies A_r .

4.4.1. *Proof Sketch.* By B_r , any ideal I as in Assumption 4.9 with $c(I) = E_n(I)$ can be written in the form

$$I = (f_{1,1}, \dots, f_{1,a_1}) + ((f'_{2,1} + f''_{2,1}) + \dots + (f'_{r,a_r} + f''_{r,a_r}))$$

where $f'_{i,j} \in k[\mathbf{x}_2, \dots, \mathbf{x}_r]$ and $f''_{i,j} \in (\mathbf{x}_1)$, and $f_{1,1}, \dots, f_{1,a_1}, f'_{2,1}, \dots, f'_{r,a_r}$ are all contained in \mathfrak{D} and form a regular sequence. If $f''_{i,j} \in \mathfrak{D}$ for all i, j , then $I \subseteq \mathfrak{D}$ and Proposition 2.39 implies that $\bar{I} = \mathfrak{D}$ as desired.

To handle the case that $I \not\subseteq \mathfrak{D}$, we suppose that I is as close to \mathfrak{D} as possible without having $I \subseteq \mathfrak{D}$. Specifically, we consider an ideal I with a unique index $2 \leq m \leq r-1$ such that $f''_{i,j} = 0$ for all $i \neq m$ and $f'_{i,j} = x_{i,j}^{d_i}$. We also suppose that for this fixed index m , the $f''_{m,j}$ are not all zero and no monomial summand of $f''_{m,j}$ is contained in \mathfrak{D} . Because of the (relatively) uncomplicated structure of I , we are able to use positive-characteristic arguments to directly show $c(I) > E_n(I)$ (Lemma 4.29).

In the general case (Lemma 4.28), we prove that any complete intersection satisfying Equation (†) but with $I \not\subseteq \mathfrak{D}$ can be degenerated to an ideal I' which is as described in the previous paragraph: close to \mathfrak{D} while satisfying $I' \not\subseteq \mathfrak{D}$. By Proposition 2.22, we have $c(I) \geq c(I') > E_n(I') = E_n(I)$.

4.4.2. Constructing the Degeneration Order.

Notation 4.18. Let $\mathbf{b}_1, \dots, \mathbf{b}_n$ denote the standard basis vectors for \mathbb{Z}^n . For $\mathbf{u} = (u_1, \dots, u_n) \in (\mathbb{Z}^+)^n$, set $|\mathbf{u}| = u_1 + \dots + u_n$. For $\mathbf{d} = (d_1, \dots, d_n) \in \mathbb{Z}^n$, we also define $\nu_{\mathbf{d}} : \mathbb{Z}^n \rightarrow \mathbb{Q}$ by $\nu_{\mathbf{d}}(\mathbf{u}) = \frac{u_1}{d_1} + \dots + \frac{u_n}{d_n}$.

Lemma 4.19. Let $0 < d_1 < \dots < d_n \in (\mathbb{Z}^+)^n$ and set $\mathbf{d} = (d_1, \dots, d_n)$. Let $S \subseteq (\mathbb{Z}_{\geq 0})^n$. Write $S = S_2 \sqcup \dots \sqcup S_{n-1}$. Assume that S satisfies the following:

- (1) For all $\mathbf{u} \in S_i$, we have $|\mathbf{u}| = d_i$.
- (2) For all $\mathbf{u} \in S$, either $u_1 \geq 1$ or $\nu_{\mathbf{d}}(\mathbf{u}) \geq 1$.
- (3) For all $2 \leq i \leq n-1$, we have $d_i \mathbf{b}_i \notin S_i$.
- (4) There exists $\mathbf{u} \in S$ with $\nu_{\mathbf{d}}(\mathbf{u}) < 1$ (and therefore $u_1 \geq 1$).

Then there exists a linear map $\lambda : \mathbb{Z}^n \rightarrow \mathbb{Z}$ and a unique index $2 \leq m \leq n-1$ such that

(A.i) For all $2 \leq i \leq n-1, i \neq m, \mathbf{u} \in S_i$ we have $\lambda(\mathbf{u}) < \lambda(d_m \mathbf{b}_m)$.

(A.ii) We have

$$(21) \quad \max_{\mathbf{u} \in S_m} \lambda(\mathbf{u}) = \lambda(d_m \mathbf{b}_m).$$

(A.iii) For any \mathbf{u} achieving the maximum in Equation (21) we have $\nu_{\mathbf{d}}(\mathbf{u}) < 1$ and $u_1 \geq 1$.

To prove Lemma 4.19, we construct a sequence of auxiliary linear maps $\lambda_0, \dots, \lambda_s : \mathbb{Z}^n \rightarrow \mathbb{Q}$.

Lemma 4.20. Assume the setup of Lemma 4.19. There exists a linear map $\lambda_0 : \mathbb{Z}^n \rightarrow \mathbb{Q}$ such that:

- (B.i) For all $2 \leq i \leq n-1, \mathbf{u} \in S_i$ we have $\lambda_0(\mathbf{u}) \leq \lambda_0(d_i \mathbf{b}_i)$.
- (B.ii) For $\mathbf{u} \in S_i$, if $\lambda_0(\mathbf{u}) = \lambda_0(d_i \mathbf{b}_i)$ then $\nu_{\mathbf{d}}(\mathbf{u}) \leq 1$.
- (B.iii) For $\mathbf{u} \in S_i$, if $\lambda_0(\mathbf{u}) = \lambda_0(d_i \mathbf{b}_i)$ and $u_1 \geq 1$, then $\nu_{\mathbf{d}}(\mathbf{u}) < 1$.
- (B.iv) There exists an index $2 \leq j \leq n-1$ and some $\mathbf{u} \in S_j$ such that $u_1 \geq 1$ and $\lambda_0(\mathbf{u}) = d_j \lambda_0(\mathbf{b}_j)$.

Proof. To start, we set

$$(22) \quad t_0 := \min_{\mathbf{u} \in S \text{ with } u_1 \geq 1} \frac{\nu_{\mathbf{d}}(\mathbf{u}) - 1}{u_1}.$$

We define $\lambda_0 : \mathbb{Z}^n \rightarrow \mathbb{Q}$ by

$$\lambda_0(\mathbf{u}) = t_0 u_1 - \nu_{\mathbf{d}}(\mathbf{u}).$$

We claim that λ_0 satisfies (B.i)-(B.iv).

(B.i) Let $2 \leq i \leq n-1, \mathbf{u} \in S_i$. Our proof that $\lambda_0(\mathbf{u}) \leq \lambda_0(d_i \mathbf{b}_i)$ splits into two cases.

$u_1 = 0$: By assumption (2) of Lemma 4.19 we have $\lambda_0(\mathbf{u}) = -\nu_{\mathbf{d}}(\mathbf{u}) \leq -1$.

$u_1 \geq 1$: We have $\lambda_0(\mathbf{u}) \leq \left(\frac{\nu_{\mathbf{d}}(\mathbf{u})-1}{u_1}\right) u_1 - \nu_{\mathbf{d}}(\mathbf{u}) = -1$. The property (B.i) then follows from the fact that $\lambda_0(d_i \mathbf{b}_i) = \nu_{\mathbf{d}}(d_i \mathbf{b}_i) = -1$ for all $2 \leq i \leq n-1$.

(B.ii) First, we observe that if $\lambda_0(\mathbf{u}) = \lambda_0(d_i \mathbf{b}_i) = -1$, then

$$(23) \quad \nu_{\mathbf{d}}(\mathbf{u}) = t_0 u_1 - \lambda_0(\mathbf{u}) = 1 + t_0 u_1.$$

By assumption (4) of Lemma 4.19, there exists $\mathbf{u} \in S$ with $u_1 \geq 1$ and $\nu_{\mathbf{d}}(\mathbf{u}) < 1$. It follows that the minimum in Equation (22) is negative, proving (B.ii).

(B.iii) The claim (B.iii) follows from Equation (23) together with the fact that $t_0 u_1 < 0$.

(B.iv) If $\mathbf{u} \in S_j$ realizes the minimum in Equation (22), then

$$\lambda_0(\mathbf{u}) = \left(\frac{\nu_{\mathbf{d}}(\mathbf{u}) - 1}{u_1} \right) u_1 + \nu_{\mathbf{d}}(\mathbf{u}) = -1 = \lambda_0(d_j \mathbf{b}_j).$$

□

Notation 4.21. Assume the notation of Lemma 4.19, and let λ_0 be as in Lemma 4.20. For $2 \leq i \leq n-1$, we define

$$S_i^{(0)} = \{\mathbf{u} \in S_i : \lambda_0(\mathbf{u}) = \lambda_0(d_i \mathbf{b}_i)\}$$

and we set $S^{(0)} = S_2^{(0)} \sqcup \cdots \sqcup S_{n-1}^{(0)}$.

By Lemma 4.20, we have some control over $S^{(0)}$.

- By definition, we have $S^{(0)} \subseteq S$.
- By (B.ii) and (B.iii), for every $\mathbf{u} \in S^{(0)}$ we have $\nu_{\mathbf{d}}(\mathbf{u}) \leq 1$ with equality only if $u_1 = 0$.
- By (B.iv), $S^{(0)}$ is nonempty. In particular, $S^{(0)}$ contains an element \mathbf{u} with $u_1 \geq 1$.

Lemma 4.22. Assume the notation of Notation 4.21. There exists a linear map $\lambda_1 : \mathbb{Z}^n \rightarrow \mathbb{Q}$ such that:

- (C.i)** For all $2 \leq i \leq n-1$, $\mathbf{u} \in S_i^{(0)}$, if $u_1 = 0$ then $\lambda_1(\mathbf{u}) < \lambda_1(d_i \mathbf{b}_i)$.
- (C.ii)** For all $2 \leq i \leq n-1$, $\mathbf{u} \in S_i^{(0)}$, if $u_1 \geq 1$ then we have $\lambda_1(\mathbf{u}) \leq \lambda_1(d_i \mathbf{b}_i)$.
- (C.iii)** There exists an index $2 \leq j \leq n-1$ and some $\mathbf{u} \in S_j^{(0)}$ such that $u_1 \geq 1$ and $\lambda_1(\mathbf{u}) = \lambda_1(d_j \mathbf{b}_j)$.

Proof. We set

$$(24) \quad t_1 := \min_{2 \leq i \leq n-1} \min_{\substack{\mathbf{u} \in S_i^{(0)} \\ \text{with } u_1 \neq 0}} \frac{d_n^2 u_2 + \cdots + d_n^m u_n - d_i d_n^i}{u_1}, \quad \lambda_1(\mathbf{u}) := t_1 u_1 - d_n^2 u_2 - \cdots - d_n^m u_n.$$

We claim that λ_1 satisfies properties (C.i)-(C.iii).

(C.i) By assumption (2) and (B.ii), we have $\nu_{\mathbf{d}}(\mathbf{u}) \leq 1$. Consequently, we deduce

$$(25) \quad 1 \geq \nu_{\mathbf{d}}(\mathbf{u}) = \frac{u_1}{d_1} + \cdots + \frac{u_i}{d_i} \geq \frac{u_1 + \cdots + u_i}{d_i}.$$

If $u_1 + \cdots + u_i = d_i$, then each inequality in Equation (25) is an equality and hence $\mathbf{u} = d_i \mathbf{b}_i$. This would violate assumption (3), so there must instead be some index $i < j \leq n$ such that $u_j \geq 1$. We then have

$$\lambda_1(\mathbf{u}) = -d_n^2 u_2 - \cdots - d_n^n u_n \leq -d_n^j < -d_i d_n^i = \lambda_1(d_i \mathbf{b}_i).$$

(C.ii) We compute

(26)

$$\lambda_1(\mathbf{u}) \leq \left(\frac{d_n^2 u_2 + \cdots + d_n^n u_n - d_i d_n^i}{u_i} \right) u_i - d_n^2 u_2 - \cdots - d_n^n u_n = -d_i d_n^i = \lambda_1(d_i \mathbf{b}_i).$$

(C.iii) By (B.iv), the set $\{\mathbf{u} \in S^{(0)} : u_1 \geq 1\}$ is nonempty. If \mathbf{u} achieves the minimum in Equation (24), then the inequality in Equation (26) is an equality. \square

Notation 4.23 (Subsuming Notation 4.21). *Let $k \geq 0$. Suppose that $\lambda_0, \dots, \lambda_k$ have been defined. For $1 \leq \ell \leq k, 2 \leq i \leq n-1$, we define*

$$S_i^{(\ell)} = \{\mathbf{u} \in S_i^{\ell-1} : \lambda_\ell(\mathbf{u}) = d_i \lambda_\ell(\mathbf{b}_i)\}$$

and set $S^{(\ell)} = S_2^{(\ell)} \sqcup \cdots \sqcup S_{n-1}^{(\ell)}$. Finally, we set $\Lambda_\ell = \{2 \leq i \leq n-1 : S_i^{(\ell)} \neq \emptyset\}$.

Similarly to the case of $S^{(0)}$, Lemmas 4.22 and 4.24 give us control over $S^{(\ell)}$ for $\ell \geq 1$. Assume we have constructed λ_{k+1} and proven Lemma 4.24 for $k \leq \ell-1$.

- By definition, we have $S^{(\ell)} \subseteq S^{(\ell-1)}$.
- By (B.iii), (C.i), and (C.ii), for all $\mathbf{u} \in S^{(\ell)}$ we have $u_1 \geq 1$ and $\nu_{\mathbf{d}}(\mathbf{u}) < 1$.
- By (C.iii) if $\ell = 1$ and (D.iii) if $\ell \geq 2$, there exists an element $\mathbf{u} \in S^{(\ell)}$ with $u_1 \geq 1$.

Lemma 4.24. *Let $k \geq 0$ and suppose that $\lambda_0, \dots, \lambda_k$ have been defined. Assume the notation of Notation 4.23 and suppose that $|\Lambda_k| \geq 2$. Set $i_k = \min \Lambda_k$. There exists $\lambda_{k+1} : \mathbb{Z}^n \rightarrow \mathbb{Q}$ such that:*

- (D.i) For $\mathbf{u} \in S_{i_k}^{(k)}$, we have $\lambda_{k+1}(\mathbf{u}) < \lambda_{k+1}(d_{i_k} \mathbf{b}_{i_k})$.
- (D.ii) For $j > i_k, \mathbf{u} \in S_j^{(k)}$, we have $\lambda_{k+1}(\mathbf{u}) \leq \lambda_{k+1}(d_j \mathbf{b}_j)$.
- (D.iii) There exists some index $j > i_k$ and some $\mathbf{u} \in S_j^{(k)}$ such that $\lambda_{k+1}(\mathbf{u}) = \lambda_{k+1}(d_j \mathbf{b}_j)$.

Proof. We define

$$(27) \quad t_{k+1} := \max_{j > i_k, \mathbf{u} \in S_j^{(k)}} \frac{u_{i_k}}{u_1}, \quad \lambda_{k+1}(\mathbf{u}) := -t_{k+1} u_1 + u_{i_k}.$$

We claim that λ_{k+1} satisfies (D.i)-(D.iii).

(D.i) For $\mathbf{u} \in S_{i_k}^{(k)}$, since $\mathbf{u} \neq d_{i_k} \mathbf{b}_{i_k}$, we have

$$\lambda_{k+1}(\mathbf{u}) \leq u_{i_k} < d_{i_k} = d_{i_k} \lambda_{k+1}(\mathbf{b}_{i_k}).$$

(D.ii) For $j > k$, $\mathbf{u} \in S_j^{(k)}$ we have

$$(28) \quad \lambda_{k+1}(\mathbf{u}) \leq -\left(\frac{u_{i_k}}{u_1}\right)u_1 + u_{i_k} = 0 = \lambda_{k+1}(d_j \mathbf{b}_j).$$

(D.iii) If \mathbf{u} attains the maximum in Equation (27), then the inequality in Equation (28) is sharp. □

Lemma 4.25 (c.f. [Eis95], Exercise 15.12). *Let $\mu_0, \dots, \mu_s : \mathbb{Z}^n \rightarrow \mathbb{Q}$ be linear maps and $U \subseteq \mathbb{Z}^n$ a finite set. There exists a map $\mu : \mathbb{Z}^n \rightarrow \mathbb{Z}$ such that for all $\mathbf{u}, \mathbf{v} \in U$, we have $\mu(\mathbf{u}) < \mu(\mathbf{v})$ if and only if*

$$(29) \quad \begin{aligned} & \text{there exists } 0 \leq k \leq s \text{ such that } \mu_k(\mathbf{u}) < \mu_k(\mathbf{v}) \\ & \text{and } \mu_\ell(\mathbf{u}) = \mu_\ell(\mathbf{v}) \text{ for all } 0 \leq \ell \leq k-1 \end{aligned}$$

We may now prove Lemma 4.19.

Proof. First, we construct the auxiliary maps $\lambda_0, \dots, \lambda_s$. Apply Lemmas 4.20 and 4.22 to S to produce linear maps λ_0, λ_1 . By (B.iv) and (C.iii), we have $|\Lambda_1| \geq 1$. We continue inductively: suppose we have constructed $\lambda_0, \dots, \lambda_k$. If $|\Lambda_k| = 1$, we set $s = k$ and proceed to the start of the following paragraph for the next step of the argument. Otherwise, apply Lemma 4.24 to produce a map λ_{k+1} . By (D.i) we have $\Lambda_{k+1} \subseteq \Lambda_k \setminus \{i_k\}$ and by (D.iii) we have $|\Lambda_{k+1}| \geq 1$. As $n-2 \geq |\Lambda_1| > |\Lambda_2| > \dots \geq 1$, we eventually arrive at an index s such that $|\Lambda_s| = 1$.

Apply Lemma 4.25 to the sequence $\lambda_0, \dots, \lambda_s$ to produce a map $\lambda : \mathbb{Z}^n \rightarrow \mathbb{Z}$. We show that λ satisfies (A.i) and (A.ii), where the index m is the unique element of Λ_s . By Lemmas 4.20, 4.22 and 4.24, for all $2 \leq i \leq n-1$, $\mathbf{u} \in S_i$, $0 \leq k \leq s$ we have $\lambda_k(\mathbf{u}) \leq \lambda_k(d_i \mathbf{b}_i)$. It follows that $\lambda(\mathbf{u}) \leq \lambda(d_i \mathbf{b}_i)$. If $\lambda(\mathbf{u}) = \lambda(d_i \mathbf{b}_i)$, then we must have $\lambda_k(\mathbf{u}) = \lambda_k(d_i \mathbf{b}_i)$ for all $0 \leq k \leq s$. Each successive equality $\lambda_k(\mathbf{u}) = \lambda_k(d_i \mathbf{b}_i)$ creates further restrictions on \mathbf{u} :

- (a) As $\lambda_0(\mathbf{u}) = \lambda_0(d_i \mathbf{b}_i)$, by Lemma 4.20 (B.ii) and (B.iii) we have either $u_1 = 0$ and $\nu_{\mathbf{d}}(\mathbf{u}) \leq 1$ or $u_1 \geq 1$ and $\nu_{\mathbf{d}}(\mathbf{u}) < 1$.
- (b) As $\lambda_1(\mathbf{u}) = \lambda_1(d_i \mathbf{b}_i)$, by Lemma 4.22 (C.i), we must in fact have $u_1 \geq 1$ and $\nu_{\mathbf{d}}(\mathbf{u}) < 1$.
- (c) As $\lambda_k(\mathbf{u}) = \lambda_k(d_i \mathbf{b}_i)$ for all $1 \leq i \leq s$, we have $i \in \Lambda_1 \cap \dots \cap \Lambda_s = \{m\}$.

Point (c) in the above list implies (A.i) and points (a), (b) imply (A.iii). For (A.ii), we have by construction that $\Lambda_s = \{m\}$ is nonempty, so there exists $\mathbf{u}^* \in S_m^{(s)} \subseteq S_m$. By Notation 4.23, the set $S_m^{(s)}$ is inductively defined as the set of $\mathbf{u} \in S_m$ such that $\lambda_k(\mathbf{u}) = \lambda_k(d_m \mathbf{b}_m)$ for all $0 \leq k \leq s$, so we have $\lambda(\mathbf{u}^*) = \lambda(d_m \mathbf{b}_m)$, hence Equation (21) holds. □

4.4.3. Applying the Degeneration Order.

Lemma 4.26. *Let R be as in Assumption 4.9. Suppose that f_1, \dots, f_{a_i} are d_i -forms comprising a regular sequence in R , and suppose that $f_j \in k[\mathbf{x}_i]$ for all $1 \leq j \leq a_i$. Then the integral closure J of (f_1, \dots, f_{a_i}) in R is equal to $(\mathbf{x}_i)^{d_i}$.*

Proof. Since $k[\mathbf{x}_i] \rightarrow R$ is faithfully flat, f_1, \dots, f_{a_i} form a regular sequence in $k[\mathbf{x}_i]$. By Theorem 2.27, the integral closure of (f_1, \dots, f_{a_i}) in $k[\mathbf{x}_i]$ is $(\mathbf{x}_i)^{d_i}$. By Proposition 2.26 (vii), we have $(\mathbf{x}_i)^{d_i} \subseteq J$. On the other hand, we have $(f_1, \dots, f_{a_i}) \subseteq (\mathbf{x}_i)^{d_i}$. By [HS06, Proposition 1.4.6], $(\mathbf{x}_i)^{d_i}$ is integrally closed in R , so $J = (\mathbf{x}_i)^{d_i}$. \square

Lemma 4.27. *Assume the setting of Assumption 4.9. Suppose I satisfies Equation (\dagger) . Then we have the following:*

- (a) *For all $1 \leq i \leq r, 1 \leq j \leq a_i$, there exists $f'_{i,j} \in k[\mathbf{x}_i]$ and $f''_{i,j} \in (\mathbf{x}_1, \dots, \mathbf{x}_{i-1})$ such that $f_{i,j} = f'_{i,j} + f''_{i,j}$.*
- (b) *For all $1 \leq i \leq r$, we have $\sqrt{I_1 + \dots + I_i} = (\mathbf{x}_1, \dots, \mathbf{x}_i)$.*
- (c) *For all $1 \leq i \leq r$, the elements $f'_{i,1}, \dots, f'_{i,a_i}$ form a regular sequence.*

Proof. The key to this proof is the following simple observation. Let $g = x_{1,1}^{e_{1,1}} \dots x_{r,a_r}^{e_{r,a_r}}$ be a monomial of degree d_i such that $w_d(g) \geq 1$. If $g \notin (\mathbf{x}_1, \dots, \mathbf{x}_{i-1})$, then we have

$$(30) \quad 1 \leq w_d(g) \leq \frac{e_{i,1} + \dots + e_{i,a_i}}{d_i} + \frac{e_{i+1,1} + \dots + e_{r,a_r}}{d_{i+1}} \\ = \frac{e_{i,1} + \dots + e_{r,a_r}}{d_i} + (d_{i+1} - d_i) \left(\frac{e_{i+1,1} + \dots + e_{r,a_r}}{d_i d_{i+1}} \right)$$

As $e_{i,1} + \dots + e_{r,a_r} = d_i$ and $\deg(g) = d_i$ we have $e_{k,\ell} = 0$ for all $k < i$. At the same time, Equation (30) implies that $e_{i+1,1} + \dots + e_{r,a_r} = 0$, so $g \in k[\mathbf{x}_i]$.

For any $1 \leq i \leq r, 1 \leq j \leq a_i$, by Equation (\dagger) we have $\text{supp}(f_{i,j}) \subseteq \mathfrak{D} \sqcup (\mathbf{x}_1)$. By the previous paragraph, we have $\text{supp}(f_{i,j}) \cap \mathfrak{D} \subseteq k[\mathbf{x}_i] \sqcup (\mathbf{x}_1, \dots, \mathbf{x}_{i-1})$. Writing $f_{i,j} = \sum_{y \in \text{supp}(f_{i,j})} \beta_y^{(i,j)} y$, we may therefore define

$$f'_{i,j} := \sum_{y \in \text{supp}(f_{i,j}) \cap k[\mathbf{x}_i]} \beta_y^{(i,j)} y, \quad f''_{i,j} = \sum_{y \in \text{supp}(f_{i,j}) \cap (\mathbf{x}_1, \dots, \mathbf{x}_{i-1})} \beta_y^{(i,j)} y,$$

which proves (a).

We prove (b) and (c) simultaneously by induction. By Equation (\dagger) , I_1 is a homogeneous ideal extended from $k[\mathbf{x}_1]$ and $\text{ht}(I_1) = a_1$, so (b) holds for $i = 1$. As $f_{1,j} = f'_{1,j}$ for $1 \leq j \leq a_1$, (c) also holds for $i = 1$. Suppose that (b), (c) hold for $i - 1$. Then $\sqrt{I_1 + \dots + I_{i-1}} = (\mathbf{x}_1, \dots, \mathbf{x}_{i-1})$, so we have

$$(31) \quad \sqrt{I_1 + \dots + I_i} = \sqrt{(\mathbf{x}_1, \dots, \mathbf{x}_{i-1}) + (f_{i,1}, \dots, f_{i,a_i})} \\ = \sqrt{(\mathbf{x}_1, \dots, \mathbf{x}_{i-1}) + (f'_{i,1}, \dots, f'_{i,a_i})}.$$

Taking the heights of all terms in Equation (31), we observe that the elements $f_{i,1}, \dots, f_{i-1,a_{i-1}}, f'_{i,1}, \dots, f'_{i,a_i}$ form a regular sequence. In particular, $f'_{i,1}, \dots, f'_{i,a_i}$ is a regular sequence, proving claim (c). As $\text{ht}((f'_{i,1}, \dots, f'_{i,a_i})) = a_i$ and $(f'_{i,1}, \dots, f'_{i,a_i})$ is a homogeneous ideal extended from $k[\mathbf{x}_i]$, we have $\sqrt{(f'_{i,1}, \dots, f'_{i,a_i})} = (\mathbf{x}_i)$. Applying Equation (31) again, we deduce claim (b). \square

Lemma 4.28. *Assume the setting of Assumption 4.9. Suppose I satisfies Equation (†) and $I \not\subseteq \mathfrak{D}$. Then there exists an integer $2 \leq m \leq r - 1$ and d_m -forms $h_{m,1}, \dots, h_{m,a_m}$ such that, setting*

$$J := (\mathbf{x}_1)^{d_1} + \dots + (\mathbf{x}_{m-1})^{d_{m-1}} + (h_{m,1}, \dots, h_{m,a_m}) + (\mathbf{x}_{m+1})^{d_{m+1}} + \dots + (\mathbf{x}_r)^{d_r},$$

we have

- (i) For all $1 \leq j \leq a_m$, we have $h_{m,j} = h'_{m,j} + h''_{m,j}$, where $h'_{m,j} \in k[\mathbf{x}_m]$, $h''_{m,j} \in (\mathbf{x}_1)$, and $w_d(y) < 1$ for all $y \in \text{supp}(h''_{m,j})$;
- (ii) There exists some j such that $h''_{m,j} \neq 0$;
- (iii) J is \mathfrak{m} -primary with $E_n(J) = E_n(I)$;
- (iv) $c(J) \leq c(I)$.

Proof. Let $f'_{i,j}, f''_{i,j}$ be as in Lemma 4.27. Define a semigroup homomorphism $\rho : \text{Mon}(\mathbf{x}_1, \dots, \mathbf{x}_r) \rightarrow \mathbb{Z}_{\geq 0}^r$ by $\rho(x_{i,j}) = \mathbf{b}_i$. For $2 \leq i \leq r$, we define

$$S_i := \bigcup_{j=1}^{a_i} \rho(\text{supp}(f''_{i,j})).$$

Set $S = S_2 \sqcup \dots \sqcup S_{r-1}$. We claim that S satisfies the hypotheses (1)-(4) of Lemma 4.19.

- (1) The claim follows from the fact that $f''_{i,j}$ is homogeneous of degree d_i and $\deg(y) = |\rho(y)|$ for any monomial $y \in R$.
- (2) Let $y \in \text{supp}(f''_{i,j})$ be a monomial. If $y \in (\mathbf{x}_1)$, then $\rho(y)_1 \geq 1$. Otherwise, as I satisfies Equation (†), we have $1 \leq w_d(y) = \nu_d(\rho(y))$.
- (3) By construction, if $y \in \text{supp}(f''_{i,j})$ then $y \in (\mathbf{x}_1, \dots, \mathbf{x}_{i-1})$, so $\rho(y)_1 + \dots + \rho(y)_{i-1} \geq 1$, hence $\rho(y) \neq d_i \mathbf{b}_i$.
- (4) By assumption, we have $I \not\subseteq \mathfrak{D}$, thus we have $f_{i,j} \notin \mathfrak{D}$ for some $1 \leq i \leq r, 1 \leq j \leq a_i$. By Equation (†), I_1 is extended from $k[\mathbf{x}_1]$, hence $I_1 \subseteq (\mathbf{x}_1)^{d_1} \subseteq \mathfrak{D}$. Moreover, by Lemma 3.3 we have $\mathfrak{m}^{d_r} \subseteq \mathfrak{D}$, hence $I_r \subseteq \mathfrak{D}$. We conclude that there exists some $2 \leq i \leq r - 1$ and some $1 \leq j \leq a_i$ such that $f_{i,j} \notin \mathfrak{D}$. As $f'_{i,j} \in \mathfrak{m}^{d_i} \cap k[\mathbf{x}_i] \subseteq \mathfrak{D}$ and $f'_{i,j} + f''_{i,j} = f_{i,j} \notin \mathfrak{D}$, it must be the case that $f''_{i,j} \notin \mathfrak{D}$. In particular, there exists a monomial y in the support of $f''_{i,j}$ such that $y \notin \mathfrak{D}$. Recalling that \mathfrak{D} consists precisely of the elements $f \in R$ with $w_d(f) \geq 1$, for this monomial y we have $\rho(y) \in S$ and $\nu_d(\rho(y)) = w_d(y) < 1$.

We now apply Lemma 4.19 to the set S to produce a map $\lambda : \mathbb{Z}^r \rightarrow \mathbb{Z}$ and an index $2 \leq m \leq r - 1$ such that

(A.i) For all $2 \leq i \leq r - 1, i \neq m, \mathbf{u} \in S_i$ we have $\lambda(\mathbf{u}) < \lambda(d_m \mathbf{b}_m)$.

(A.ii) We have

$$(21) \quad \max_{\mathbf{u} \in S_m} \lambda(\mathbf{u}) = \lambda(d_m \mathbf{b}_m).$$

(A.iii) For any \mathbf{u} achieving the maximum in Equation (21) we have $\nu_d(\mathbf{u}) < 1$ and $u_1 \geq 1$.

Let in_λ denote the monomial partial order given by $y <_\lambda z$ if $\lambda(\rho(y)) < \lambda(\rho(z))$. We consider the leading terms $\text{in}_\lambda(f_{i,j})$ for $2 \leq i \leq r-1, 1 \leq j \leq a_i$. For any $y \in \text{supp}(f'_{i,j})$, we have $\rho(y) = d_i \mathbf{b}_i$, and in particular $\lambda(\rho(y)) \leq \lambda(d_i \mathbf{b}_i)$. By (A.i) and (A.ii), we have $\lambda(\rho(y)) \leq \lambda(d_i \mathbf{b}_i)$ for all $y \in \text{supp}(f''_{i,j})$. It follows that

$$\max_{y \in \text{supp}(f_{i,j})} \lambda(\rho(y)) \leq \lambda(d_i \mathbf{b}_i).$$

By Lemma 4.27, the elements $f'_{i,j}$ form a regular sequence. In particular, $f'_{i,j} \neq 0$, so $\text{supp}(f_{i,j}) \cap k[\mathbf{x}_i] = \text{supp}(f'_{i,j}) \neq \emptyset$. If $y_{i,j} \in \text{supp}(f'_{i,j})$, then we have

$$\max_{y \in \text{supp}(f_{i,j})} \lambda(\rho(y)) \geq \lambda(\rho(y_{i,j})) = \lambda(d_i \mathbf{b}_i).$$

We now have an expression for the leading term $\text{in}_\lambda(f_{i,j})$. If we write $f_{i,j} = \sum_{y \in \text{supp}(f_{i,j})} \beta_y^{(i,j)} y$, then $\text{in}_\lambda(f_{i,j})$ is the sum over all $\gamma_y^{(i,j)} y$ such that $\lambda(\rho(y))$ is as large as possible. To be precise, we have

$$(32) \quad \text{in}_\lambda(f_{i,j}) = \sum_{y \in \text{supp}(f_{i,j}): \lambda(\rho(y)) = \lambda(d_i \mathbf{b}_i)} \beta_y^{(i,j)} y = f'_{i,j} + \sum_{y \in \text{supp}(f''_{i,j}): \lambda(\rho(y)) = \lambda(d_i \mathbf{b}_i)} \beta_y^{(i,j)} y.$$

For $1 \leq i \leq a_m$, set $h_{m,j} := \text{in}_\lambda(f_{m,j})$. With this choice of $h_{m,1}, \dots, h_{m,a_m}$, we verify that J satisfies conclusions (i)-(iv) of the lemma.

- (i) We set $h'_{m,j} := f'_{m,j}$ and $h''_{m,j} := \sum_{y \in \text{supp}(f''_{m,j}): \lambda(\rho(y)) = \lambda(d_m \mathbf{b}_m)} \gamma_y^{(i,j)} y$. By Equation (32), we have $h'_{m,j} + h''_{m,j} = \text{in}_\lambda(f_{m,j}) = h_{m,j}$. By the definition of $f'_{m,j}$ in Lemma 4.27, we have $\text{supp}(h'_{m,j}) \subseteq k[\mathbf{x}_m]$. For any $y \in \text{supp}(h''_{m,j})$, we have $y \in \text{supp}(f''_{m,j})$, so $\rho(y) \in S_m$. We also have $\lambda(\rho(y)) = \lambda(d_m \mathbf{b}_m)$, hence by (A.iii) we have $\rho(y)_1 \geq 1$ and $\nu_{\mathbf{d}}(\rho(y)) < 1$, hence $y \in (\mathbf{x}_1)$ and $w_{\mathbf{d}}(y) < 1$.
- (ii) By (A.ii), there exists $\mathbf{u} \in S_m$ such that $\lambda(\mathbf{u}) = \lambda(d_m \mathbf{b}_m)$. By the definition of S_m , there exists some index $1 \leq j \leq a_m$ and some $y \in \text{supp}(f''_{m,j})$ such that $\rho(y) = \mathbf{u}$. By Equation (32), we have $y \in \text{supp}(h''_{m,j})$.
- (iii) We argue similarly to Equation (31). By Lemma 4.26, we have $\sqrt{(f'_{m,1}, \dots, f'_{m,a_m})} = (\mathbf{x}_m)$. By Equation (32), we have $h_{m,j} \equiv f'_{m,j} \pmod{(\mathbf{x}_1)}$, so it follows that

$$\begin{aligned} \sqrt{J} &= \sqrt{(\mathbf{x}_1) + \dots + (\mathbf{x}_{m-1}) + (h_{m,1}, \dots, h_{m,a_m}) + (\mathbf{x}_{m+1}) + \dots + (\mathbf{x}_r)} \\ &= \sqrt{(\mathbf{x}_1) + \dots + (\mathbf{x}_{m-1}) + (f'_{m,1}, \dots, f'_{m,a_m}) + (\mathbf{x}_{m+1}) + \dots + (\mathbf{x}_r)} = \mathbf{m}. \end{aligned}$$

To see that $E_n(J) = E_n(I)$, we note that J has the same integral closure as the ideal

$$J' = (x_{1,1}^{d_1}, \dots, x_{m-1,a_{m-1}}^{d_{m-1}}, h_{m,1}, \dots, h_{m,a_m}, x_{m+1,1}^{d_{m+1}}, \dots, x_{r,a_r}^{d_r}).$$

As J' is a complete intersection, by Lemma 3.8 we have

$$E_n(J) = E_n(J') = \frac{a_1}{d_1} + \dots + \frac{a_r}{d_r} = E_n(I).$$

- (iv) Let $I' = (\text{in}_\lambda(f_{1,1}), \dots, \text{in}_\lambda(f_{r,a_r}))$. We claim that $J \subseteq \overline{I'}$. By Equation (32) and (A.iii), we have $\text{in}_\lambda(f_{i,j}) = f'_{i,j}$ for $i \neq m$. For $i \neq m$, it follows from Lemmas 4.26 and 4.27 that $(\mathbf{x}_i)^{d_i} = \overline{(f'_{i,1}, \dots, f'_{i,a_i})} \subseteq \overline{I'}$. The claim that $J \subseteq \overline{I'}$ follows once we note that $h_{m,j} = \text{in}_\lambda(f_{i,j}) \in I'$. To see that $c(J) \leq c(I)$, by Propositions 2.10 and 2.22 we have

$$c(J) = c(J') \leq c(\overline{I'}) = c(I') \leq c(\text{in}_\lambda(I)) \leq c(I).$$

□

Lemma 4.29. *Assume the setting of Assumption 4.9. Let $J \subseteq R$ be an ideal satisfying conditions (i)-(iii) of Lemma 4.28. Then $c(J) > E_n(J)$.*

Proof. Set $J_m = (h_{m,1}, \dots, h_{m,a_m})$. We first prove this result in characteristic $p > 0$.

We claim that the assumptions on J_m imply that $\text{ess}(J_m) > a_m$. To see this, suppose on the contrary that $\text{ess}(J_m) \leq a_m$. As $\text{ht}(J_m) = a_m$, it must be the case that $\text{ess}(J_m) = a_m$. By [Bai25, Lemma 3.14] we have $J_m \subseteq (\ell_1, \dots, \ell_{a_m})^d$ for some linearly independent 1-forms $\ell = \ell_1, \dots, \ell_{a_m} \in R$. In particular, $\sqrt{J_m} = (\ell_1, \dots, \ell_{a_m})$.

We can compute $\sqrt{J_m}$ from a different perspective. By assumption, for all $y \in \text{supp}(h''_{m,j})$ we have $y \in (\mathbf{x}_1) \setminus \mathfrak{D}$. In particular, $y \notin (\mathbf{x}_1)^{d_m}$, so we have instead $y \in (\mathbf{x}_1) \cap (\mathbf{x}_2, \dots, \mathbf{x}_r)$. It follows that

$$(\ell) = \sqrt{J_m} \subseteq \sqrt{(\mathbf{h}'_m) + (\mathbf{h}''_m)} \subseteq (\mathbf{x}_m) + (\mathbf{x}_1) \cap (\mathbf{x}_2, \dots, \mathbf{x}_r)$$

which implies that $(\ell) = (\mathbf{x}_m)$. This contradicts the assumption that $h''_{m,j} \neq 0$ for some index $1 \leq j \leq a_m$, so we conclude that $\text{ess}(J_m) > a_m$. By [Bai25, Theorem 3.17] we have $c(J_m) > \frac{a_m}{d_m}$.

Let $f = h_{1,m}^{t_1} \dots h_{m,a_m}^{t_{a_m}}$ be a homogeneous generator of $(J_m)^{\nu_{J_m}(p^e)}$ such that $f \notin \mathfrak{m}^{[p^e]}$. Write

$$f = \sum_{y \in \text{supp}(f)} \beta_y y, \quad f' := \sum_{y \in \text{supp}(f) \setminus \mathfrak{m}^{[p^e]}} \beta_y y.$$

As $f \equiv f' \pmod{\mathfrak{m}^{[p^e]}}$, we have $f' J_m \subseteq \mathfrak{m}^{[p^e]}$.

Applying Theorem 2.29 and Lemma 4.26 to the ideal $\frac{J_m + (\mathbf{x}_1)}{(\mathbf{x}_1)}$, we have

$$\frac{(\mathbf{x}_m)^{a_m d_m} + (\mathbf{x}_1)}{(\mathbf{x}_1)} \subseteq \frac{J_m + (\mathbf{x}_1)}{(\mathbf{x}_1)}.$$

Let μ_1, \dots, μ_M be a minimal set of monomial generators for $(\mathbf{x}_m)^{a_m d_m}$. Subsequently, we let $\widetilde{\mu}_1, \dots, \widetilde{\mu}_M \in J_m$ be homogeneous elements such that $\mu_i - \widetilde{\mu}_i \in (\mathbf{x}_1)$ for all $1 \leq i \leq M$. Let \succ denote the reverse lexicographic order after putting the variables in the order $\mathbf{x}_1 \prec \dots \prec \mathbf{x}_r$. By definition of the reverse lexicographic order, we have $y \prec z$ for any $y \in (\mathbf{x}_1), z \in k[\mathbf{x}_2, \dots, \mathbf{x}_r]$. In particular, we have $\text{in}_\succ(\widetilde{\mu}_i) = \text{in}_\succ(\mu_i + (\widetilde{\mu}_i - \mu_i)) = \mu_i$. It follows that $(\mathbf{x}_m)^{a_m d_m} \subseteq \text{in}_\succ(J_m)$.

By construction, we have $f \notin \mathfrak{m}^{[p^e]}$ and $f J_m \subseteq J_m^{\nu_{J_m}(p^e)+1} \subseteq \mathfrak{m}^{[p^e]}$. As $f' \equiv f \pmod{\mathfrak{m}^{[p^e]}}$, we have $f' \notin \mathfrak{m}^{[p^e]}$ and $f' J_m \subseteq \mathfrak{m}^{[p^e]} + J_m^{\nu_{J_m}(p^e)+1} \subseteq \mathfrak{m}^{[p^e]}$. Let $y :=$

$\text{in}_>(f')$. As no element of $\text{supp}(f')$ is in $\mathfrak{m}^{[p^e]}$, we have $y \notin \mathfrak{m}^{[p^e]}$. Additionally, we have

$$\text{in}_>(f') \text{in}_>(J_m) \subseteq \text{in}_>(f' J_m) \subseteq \text{in}_>(\mathfrak{m}^{[p^e]}) = \mathfrak{m}^{[p^e]}.$$

By Lemma 4.3, we have

$$(\mathfrak{m}^{[p^e]} : J_m) \subseteq (\mathfrak{m}^{[p^e]} : (\mathbf{x}_m)^{a_m d_m}) = \mathfrak{m}^{[p^e]} + (\mathbf{x}_m)^{a_m(p^e-1)-a_m d_m+1}.$$

As $y \notin \mathfrak{m}^{[p^e]}$, we have

$$(33) \quad \text{ord}_{(\mathbf{x}_m)}(y) \geq a_m(p^e - 1) - a_m d_m + 1.$$

Additionally, we write

$$f = \prod_{j=1}^{a_m} (h'_{m,j} + h''_{m,j})^{t_j} = \prod_{j=1}^{a_m} \sum_{\substack{0 \leq t'_j, t''_j \leq t_j \\ t'_j + t''_j = t_j}} \binom{t_j}{t'_j} (h'_{m,j})^{t'_j} (h''_{m,j})^{t''_j}.$$

As $\text{supp}(f') \subseteq \text{supp}(f)$, there exist $(t'_1, t''_1), \dots, (t'_{a_m}, t''_{a_m})$ such that $t'_j + t''_j = t_j$ for all $1 \leq j \leq a_m$ and

$$y \in \text{supp} \left((h'_{m,1})^{t'_1} (h''_{m,1})^{t''_1} \dots (h'_{m,a_m})^{t'_{a_m}} (h''_{m,a_m})^{t''_{a_m}} \right).$$

Set $J'_m = (h'_{m,1}, \dots, h'_{m,a_m})$. As J'_m is extended from $k[\mathbf{x}_m]$, by Lemma 4.26 we have $J'_m \subseteq (\mathbf{x}_m)^{d_m}$, so

$$t'_1 + \dots + t'_{a_m} \leq \nu_{J'_m}(p^e) \leq \nu_{(\mathbf{x}_m)^{d_m}}(p^e) \leq \left\lfloor \frac{a_m(p^e - 1)}{d_m} \right\rfloor.$$

By assumption, $J_m \not\subseteq \mathfrak{D}$. Define

$$\sigma := \max_{\substack{1 \leq j \leq a_m \\ y \in \text{supp}(h_{m,j}) \setminus (\mathbf{x}_m)^{d_m}}} w_d(y),$$

which satisfies $\sigma < 1$ by conditions (ii) and (iv). Consequently, we have

$$\begin{aligned} w_d(y) &\leq \sum_{j=1}^{a_m} t'_j w_d(h'_{m,j}) + \sum_{j=1}^{a_m} t''_j \left(\max_{z \in \text{supp}(h''_{m,j})} w_d(z) \right) \\ &= \sum_{j=1}^{a_m} t'_j + \sum_{j=1}^{a_m} t''_j \left(\max_{z \in \text{supp}(h''_{m,j})} w_d(z) \right) \leq \sum_{j=1}^{a_m} t'_j + \sigma \sum_{j=1}^{a_m} t''_j \\ (34) \quad &\leq \left\lfloor \frac{a_m(p^e - 1)}{d_m} \right\rfloor + \sigma \left(\nu_{J_m}(p^e) - \left\lfloor \frac{a_m(p^e - 1)}{d_m} \right\rfloor \right). \end{aligned}$$

As in Lemma 4.3, let $\mathfrak{a}_\beta, \mathfrak{a}_\beta^+$ denote the ideals $\{f \in R : w_d(f) \geq \beta\}, \{f \in R : w_d(f) > \beta\}$ respectively. Let t_e denote the quantity in Equation (34); as $w_d(y) \leq t_e$ we have $y \notin \mathfrak{a}_{t_e}^+$. By assumption, $y \notin \mathfrak{m}^{[p^e]}$, and as y is a monomial we have $y \notin \mathfrak{m}^{[p^e]} + \mathfrak{a}_{t_e}^+$. Set $u_e := (p^e - 1) \left(\frac{a_1}{d_1} + \dots + \frac{a_r}{d_r} \right)$. It follows from Lemma 4.3 that

$$y \notin \mathfrak{m}^{[p^e]} + \mathfrak{a}_{t_e}^+ = (\mathfrak{m}^{[p^e]} : \mathfrak{a}_{u_e - t_e}).$$

Let $z \in \mathfrak{a}_{u_e - t_e}$ such that $yz \notin \mathfrak{m}^{[p^e]}$. Write $z = z'z''$ where $z'' \in k[\mathbf{x}_m]$ and $z' \in k[\mathbf{x}_1, \dots, \widehat{\mathbf{x}}_m, \dots, \mathbf{x}_r]$. As $yz \notin \mathfrak{m}^{[p^e]}$, by Equation (33) we have

$$\text{ord}_{(\mathbf{x}_m)}(z'') \leq (p^e - 1)a_m - \text{ord}_{(\mathbf{x}_m)}(y) \leq a_m d_m,$$

hence $w_d(z') = w_d(z) - w_d(z'') \geq u_e - t_e - a_m$. As $\mathfrak{D} = \overline{(\mathbf{x}_1)^{d_1} + \dots + (\mathbf{x}_r)^{d_r}}$, by Theorem 2.29 we have

$$z' \in \mathfrak{D}^{\lfloor u_e - t_e \rfloor - a_m} \subseteq ((\mathbf{x}_1)^{d_1} + \dots + (\mathbf{x}_r)^{d_r})^{\lfloor u_e - t_e \rfloor - a_m - n}.$$

Since $z' \notin (\mathbf{x}_m)$, we in fact have

$$z' \in ((\mathbf{x}_1)^{d_1} + \dots + (\mathbf{x}_{m-1})^{d_{m-1}} + (\mathbf{x}_{m+1})^{d_{m+1}} + \dots + (\mathbf{x}_r)^{d_r})^{\lfloor u_e - t_e \rfloor - a_m - n} \subseteq (J)^{\lfloor u_e - t_e \rfloor - a_m - n}.$$

As $z'f \notin \mathfrak{m}^{[p^e]}$, it follows that $\nu_J(p^e) \geq \nu_{J_m}(p^e) + \lfloor u_e - t_e \rfloor - a_m - n$. Dividing by p^e and letting $e \rightarrow \infty$, we obtain

$$\begin{aligned} c(J) &\geq c(J_m) + \lim_{e \rightarrow \infty} \frac{u_e}{p^e} - \lim_{e \rightarrow \infty} \frac{t_e}{p^e} \\ &= c(J_m) + \left(\frac{a_1}{d_1} + \dots + \frac{a_r}{d_r} \right) - \left(\frac{a_m}{d_m} (1 - \sigma) + \sigma c(J_m) \right) \\ &= (1 - \sigma) \left(c(J_m) - \frac{a_m}{d_m} \right) + \left(\frac{a_1}{d_1} + \dots + \frac{a_r}{d_r} \right) \end{aligned}$$

Since $\sigma < 1$ and $c(J_m) > \frac{a_m}{d_m}$, it follows that the above quantity exceeds $E_n(J)$.

In characteristic zero, one notes that for any ideal J satisfying conditions (i)-(iii), the reduction of the pair (R, J) to characteristic $p \gg 0$ satisfies conditions (i)-(iii). Moreover, the quantity σ is constant for $p \gg 0$. Assuming the reduction notation of Assumption 4.1, we have

$$\begin{aligned} c(J) &= \lim_{\substack{\mu \in \text{Spec } A \\ \text{char } A/\mu \rightarrow \infty}} c(J_\mu) \geq (1 - \sigma) \left(\lim_{\substack{\mu \in \text{Spec } A \\ \text{char } A/\mu \rightarrow \infty}} c(J_{m,\mu}) - \frac{a_m}{d_m} \right) + E_n(J) \\ &= (1 - \sigma) \left(c(J_m) - \frac{a_m}{d_m} \right) + E_n(J) > E_n(J). \end{aligned}$$

□

Lemmas 4.16, 4.28 and 4.29 combine to give us a proof of Theorem 5.1 in the case of an \mathfrak{m} -primary complete intersection.

Proposition 4.30. *Assume the setup of Assumption 4.9 and suppose $c(I) = E_n(I)$. Then there exists $\gamma \in \text{GL}_n(k)$ such that $\overline{\gamma I} = \mathfrak{D}$.*

Proof. Using Lemma 4.16, we produce $\gamma \in \text{GL}_n(k)$ such that γI satisfies Equation (†). By Lemmas 4.28 and 4.29, we have $\overline{\gamma I} = \mathfrak{D}$. □

5. PROOF OF THEOREM B

In this section, we prove the main result of this article, which we restate below.

Theorem 5.1. *Let k be a perfect field. Let $R = k[x_1, \dots, x_n]$ and let $I \subseteq R$ be a homogeneous ideal with $\text{ht}(I) \geq l$. If $E_l(I) = c(I)$, then there exist integers d_1, \dots, d_l and $\gamma \in \text{GL}_n(k)$ such that*

$$\gamma \bar{I} = \overline{(x_1^{d_1}, \dots, x_l^{d_l})}.$$

5.1. Algebraically closed fields. Throughout this subsection, fix the following assumption.

Assumption 5.2. *Let L be an uncountable, algebraically closed field. Let $S = L[x_1, \dots, x_n]$ and let $I \subseteq S$ be a homogeneous. As in Lemma 3.2, for $1 \leq j \leq n$, let $S_j := L[x_1, \dots, x_j]$, $\pi_j : S \rightarrow S_j$ the natural projection map and $\iota_j : S_j \rightarrow S$ the natural embedding. Let $>$ denote the graded reverse lexicographic order $>_{\text{grlex}}$.*

Definition 5.3. Let k be a field, $R = k[x_1, \dots, x_n]$, $I \subseteq R$ a homogeneous ideal, and $t \in \mathbb{Z}^+$. We let $[I]_t$ denote the vector space of t -forms in I and we let $[I]_{\leq t}$ denote the direct sum of the $[I]_s$ for $s \leq t$.

Definition 5.4. Let k be a field, $R = k[x_1, \dots, x_n]$, and $I \subseteq R$ a homogeneous ideal of height $l > 0$. For $1 \leq i \leq l$, let $d_i(I) = \min\{j : \text{ht}[I]_{\leq j} R \geq i\}$.

Lemma 5.5 (c.f. [May14]). *Assume the setting of Assumption 5.2. Let $\gamma \in \text{GL}_n(L)$ be very general: for now, we impose the condition that for all $m > 0$, we have $\text{in}_>(\gamma I^m) = \text{gin}_>(I^m)$; we will impose countably many additional conditions in Lemma 5.10. For $1 \leq j \leq n, m > 0$, set $\mathbf{a}_{j,m} := \text{in}_>(\pi_j(\gamma I^m)) \subseteq S_j$. For $j > 0, 1 \leq i \leq j$, let $\mathbf{b}_i^{(j)}$ denote the i th unit vector of \mathbb{R}^j . For $j \geq i$, let $p_j(i) := \inf\{t : t\mathbf{b}_i^{(j)} \in \Gamma(\mathbf{a}_{j,\bullet})\}$. Then for all j , we have $p_j(j) = p_n(j)$.*

Proof. By Lemma 3.2 we have $\iota_j(\mathbf{a}_{j,\bullet}) \subseteq \mathbf{a}_{n,\bullet}$ for all $1 \leq j \leq n$, so $p_n(j) \leq p_j(j)$. For the reverse direction, set $t = p_n(j)$. Since $t\mathbf{b}_j^{(n)} \in \bigcup_{m>0} \frac{1}{2^m} \Gamma(\mathbf{a}_{n,2^m})$, there exists a sequence $\{\mathbf{a}_m = (a_{m,1}, \dots, a_{m,n})\}_{m>0}$ such that $\mathbf{a}_m \in \Gamma(\mathbf{a}_{n,2^m})$ for all m and $\lim_{m \rightarrow \infty} 2^{-m} \mathbf{a}_m = t\mathbf{b}_j^{(n)}$. For any choice of $\{\mathbf{a}_m\}_{m>0}$, we also have $(\lceil a_{m,1} \rceil, \dots, \lceil a_{m,n} \rceil) \in \Gamma(\mathbf{a}_{n,2^m})$ and $\lim_{m \rightarrow \infty} \frac{(\lceil a_{m,1} \rceil, \dots, \lceil a_{m,n} \rceil)}{2^m} = t\mathbf{b}_j^{(n)}$. We may therefore assume without loss of generality that $\mathbf{a}_m \in \mathbb{Z}_{\geq 0}^n$ for all $m > 0, 1 \leq i \leq n$, hence for all $m > 0$, we have $\mathbf{x}^{\mathbf{a}_m} \in \overline{\mathbf{a}_{n,2^m}}$.

By [GW15, Theorem 2.1], $\overline{\mathbf{a}_{n,2^m}}$ is Borel-fixed, so

$$x_1^{a_{m,1}} \cdots x_{j-1}^{a_{m,j-1}} x_j^{a_{m,j} + \cdots + a_{m,n}} \in \overline{\mathbf{a}_{n,2^m}}.$$

We further note that $\mathbf{a}_{j,m} = \pi_j(\mathbf{a}_{n,m})$ by Lemma 3.2. By Proposition 2.26(iii), we conclude

$$\pi_j(x_1^{a_{m,1}} \cdots x_{j-1}^{a_{m,j-1}} x_j^{a_{m,j} + \cdots + a_{m,n}}) \in \pi_j(\overline{\mathbf{a}_{n,2^m}}) \subseteq \overline{\pi_j(\mathbf{a}_{n,2^m})} = \overline{\mathbf{a}_{j,2^m}}.$$

It follows that

$$t\mathbf{b}_j^{(j)} = \lim_{m \rightarrow \infty} \frac{(a_{m,1}, \dots, a_{j-1}, a_j + \cdots + a_n)}{m} \in \Gamma(\mathbf{a}_{j,\bullet}),$$

which proves $p_j(j) \leq t = p_n(j)$. \square

Remark 5.6. Assume the setup of Lemma 5.5. A priori, there exists a sequence

$$\{\mathbf{a}_m = (a_{m,1}, \dots, a_{m,j})\}_{m>0} \subseteq \mathbb{R}_{\geq 0}^j$$

such that for all $m > 0$ we have $\lim_{m \rightarrow \infty} 2^{-m} \mathbf{a}_m = t \mathbf{b}_j^{(j)}$ and $\mathbf{a}_m \in \Gamma(\mathfrak{a}_{2^m})$. Lemma 5.7 is a minor refinement, allowing us to assume that $\mathbf{x}^{\mathbf{a}_m} \in \mathfrak{a}_{2^m}$.

Lemma 5.7. *Assume the setup of Lemma 5.5. There exists a sequence $\{\mathbf{a}'_m\}_{m>0} \subseteq \mathbb{Z}^j$ such that $\mathbf{x}^{\mathbf{a}'_m} \in \mathfrak{a}_{j,2^m}$ for all $m > 0$ and $\lim_{m \rightarrow \infty} 2^{-m} \mathbf{a}'_m = p_j(j) \mathbf{b}_j^{(j)}$.*

Proof. Let $\{\mathbf{a}_m\}_m$ be as in Remark 5.6 and set $\mathbf{a}'_m = (2^{m-\lfloor m/2 \rfloor} + j - 1) \mathbf{a}_{\lfloor m/2 \rfloor}$. By Theorem 2.29, we have

$$\mathbf{x}^{\mathbf{a}'_m} = (\mathbf{x}^{\mathbf{a}_{\lfloor m/2 \rfloor}})^{2^{m-\lfloor m/2 \rfloor} + j - 1} \in \overline{\mathfrak{a}_{2^{\lfloor m/2 \rfloor}}^{2^{m-\lfloor m/2 \rfloor} + j - 1}} \subseteq \mathfrak{a}_{2^{\lfloor m/2 \rfloor}}^{2^{m-\lfloor m/2 \rfloor}} \subseteq \mathfrak{a}_{2^m}.$$

The result follows once we note that

$$\lim_{m \rightarrow \infty} 2^{-m} \mathbf{a}'_m = \lim_{m \rightarrow \infty} \frac{2^{m-\lfloor m/2 \rfloor} + j - 1}{2^{m-\lfloor m/2 \rfloor}} 2^{-\lfloor m/2 \rfloor} \mathbf{a}_{\lfloor m/2 \rfloor} = p_j(j) \mathbf{b}_j^{(j)}.$$

\square

Lemma 5.8. *Let k be an algebraically closed field and $R = k[x_1, \dots, x_j]$. Let \mathfrak{q} be a homogeneous prime ideal of height $j - 1$ with $x_j \notin \mathfrak{q}$. If $>$ denotes the reverse lexicographic order, then for all $m > 0$ we have $\text{in}_{>}(\mathfrak{q}^m) = (x_1, \dots, x_{j-1})^m$.*

Proof. Since k is algebraically closed, there exist linear forms $\ell_1, \dots, \ell_{j-1} \in R_1$ such that $\mathfrak{q} = (\ell_1, \dots, \ell_{j-1})$. By [Eis95, Theorem 15.17], \mathfrak{q} and $\text{in}_{>}(\mathfrak{q})$ have the same Hilbert series. Moreover, $\text{in}_{>}(\mathfrak{q})$ is a monomial ideal not containing x_j , so we must have $\text{in}_{>}(\mathfrak{q}) = (x_1, \dots, x_{j-1})$. For $m > 1$, we have the standard containment $(x_1, \dots, x_{j-1})^m = \text{in}_{>}(\mathfrak{q})^m \subseteq \text{in}_{>}(\mathfrak{q}^m)$. As $(x_1, \dots, x_{j-1})^m$ has the same Hilbert series as \mathfrak{q}^m , the result follows. \square

Lemma 5.9. *Let k be a field, $R = k[x_1, \dots, x_j]$, and $\mathfrak{n} = (x_1, \dots, x_j)$. If $I \subseteq R$ is a homogeneous ideal, then*

$$[I\mathfrak{n}^t]_u = [[I]_{u-t}R]_u.$$

Proof. First, we verify the following: if f is homogeneous of degree $u - t$, then $[f\mathfrak{n}^t]_u = [fR]_u$. The containment \subseteq is clear. For \supseteq , if g is homogeneous and $gf \in [fR]_u$, then $g \in \mathfrak{n}^t$, so $gf \in [f\mathfrak{n}^t]_u$.

Now let f_1, \dots, f_r be homogeneous elements spanning $[I]_{u-t}$, and let g_1, \dots, g_s be homogeneous elements such that $f_1, \dots, f_r, g_1, \dots, g_s$ generate I and $\deg(g_i) > u - t$. By the first paragraph, we compute

$$[I\mathfrak{n}^t]_u = [(f_1, \dots, f_r)\mathfrak{n}^t]_u + [(g_1, \dots, g_s)\mathfrak{n}^t]_u = \sum_{i=1}^r [f_i\mathfrak{n}^t]_u = \sum_{i=1}^r [f_i]_u = [[I]_{u-t}R]_u.$$

\square

Lemma 5.10. *Assume the setup of Lemmas 5.5 and 5.7. Then for all $1 \leq j \leq n$, we have $p_n(j) = d_j(I)$.*

Proof. Set $d_j := d_j(I)$. Before we begin the proof, we first state the additional generality conditions on γ . For all $m > 0$, assume that $\text{in}_>(\pi_j(\gamma I^m)) = \text{gin}_>(\pi_j(\gamma I^m))$. Since $\text{ht } \pi_j(\gamma[I]_{\leq d_j-1}R) \leq \text{ht } \gamma[I]_{\leq d_j-1}R < j$, there is some $1 \leq i \leq j$ such that $x_i \notin \sqrt{\pi_j(\gamma[I]_{\leq d_j-1}R)}$; we choose γ such that $x_j \notin \sqrt{\pi_j(\gamma[I]_{\leq d_j-1}R)}$. Each of these conditions is satisfied by a general choice of γ , so they may be realized simultaneously by a very general choice of γ .

Set $J = \pi_j(\gamma I)$. Write $J_1 = [J]_{\leq d_j-1}R$, and let J_2 be the ideal generated by the minimal homogeneous generators of J with degree at least d_j so that $J = J_1 + J_2$ and $J_2 \subseteq \mathfrak{m}^{d_j}$. By construction of γ , in the language of Lemma 5.5 we have $\mathfrak{a}_{j,m} = \text{in}_>(J^m)$. Also by construction, we have $x_j \notin \sqrt{[J]_{\leq d_j-1}}$, so we may choose a minimal prime \mathfrak{p} over J_1 such that $x_j \notin \mathfrak{p}$. As $\text{ht } \mathfrak{p} \leq j-1$, we may choose a homogeneous prime ideal $\mathfrak{q} \supseteq \mathfrak{p}$ such that $\text{ht } \mathfrak{q} = j-1$ and $x_j \notin \mathfrak{q}$. By Lemma 5.8, we have $\text{in}_>(\mathfrak{q}^m) = (x_1, \dots, x_{j-1})^m$ for all $m > 0$.

By Lemma 5.7, choose a sequence $\{\mathfrak{a}_m\}_{m>0}$ such that $\mathbf{x}^{\mathfrak{a}_m} \in \mathfrak{a}_{2^m}$ for all $m > 0$ and $\lim_{m \rightarrow \infty} 2^{-m} \mathfrak{a}_m = p_j(j) \mathbf{b}_j^{(j)}$. Let $e_m := a_{m,1} + \dots + a_{m,j}$. For all $m > 0$, we have

$$(35) \quad [J^{2^m}]_{e_m} = \left[\sum_{\substack{a+b=2^m \\ a+bd_j \leq e_m}} J_1 + J_2 \right]_{e_m}$$

We have shown that $J_1 \subseteq \mathfrak{q}$ and $J_2 \subseteq \mathfrak{m}^{d_j}$. By Equation (35) and Lemma 5.9, we have

$$\begin{aligned} \mathbf{x}^{\mathfrak{a}_m} \in [J^{2^m}]_{e_m} &\subseteq \left[\sum_{\substack{a+b=2^m \\ a+bd_j \leq e_m}} \mathfrak{q}^a \mathfrak{m}^{bd_j} \right]_{e_m} = \sum_{\substack{a+b=2^m \\ a+bd_j \leq e_m}} [\mathfrak{q}^a \mathfrak{m}^{bd_j}]_{e_m} \\ &= \sum_{\substack{a+b=2^m \\ a+bd_j \leq e_m}} = \left[\mathfrak{q}^{2^m - \left\lfloor \frac{e_m}{d_j} \right\rfloor} \right]_{e_m}. \end{aligned}$$

Taking initial terms of both sides, we have $\mathbf{x}^{\mathfrak{a}_m} \in (x_1, \dots, x_{j-1})^{2^m - \left\lfloor \frac{e_m}{d_j} \right\rfloor}$. Consequently, we have $a_{m,1} + \dots + a_{m,j-1} \geq 2^m - \left\lfloor \frac{e_m}{d_j} \right\rfloor$. As $\lim_{m \rightarrow \infty} 2^{-m}(a_{m,1} + \dots + a_{m,j-1}) = 0$, this yields

$$0 \geq \liminf_{m \rightarrow \infty} 2^{-m} \left(2^m - \left\lfloor \frac{e_m}{d_j} \right\rfloor \right) = 1 - \frac{1}{d_j} \limsup_{m \rightarrow \infty} \frac{a_{m,j}}{2^m} = 1 - \frac{p_j(j)}{d_j}.$$

From the above equation, we have $p_j(j) \geq d_j$. For the reverse containment, we have by Lemma 3.3 that $\mathfrak{m}^{d_j} \subseteq \bar{J}$. It follows from Theorem 2.29 that $x_j^{(m+j-1)d_j} \in J^m$ for all $m > 0$, hence $p_j(j) \leq d_j$. \square

Lemma 5.11. *Let $R = k[x_1, \dots, x_n]$ and let $\mathfrak{m} = (x_1, \dots, x_n)$. Let $I \subseteq R$ be a homogeneous ideal and $J \subseteq \mathfrak{m}$ any ideal. Then we have*

$$\bigcap_{t>0} \overline{I + J^t} = \bar{I}.$$

Proof. By [HS06, Corollary 6.8.5], we have

$$\overline{IR_{\mathfrak{m}}} \subseteq \bigcap_{t>0} \overline{IR_{\mathfrak{m}} + J^t R_{\mathfrak{m}}} \subseteq \bigcap_{t>0} \overline{IR_{\mathfrak{m}} + \mathfrak{m}^t R_{\mathfrak{m}}} = \overline{IR_{\mathfrak{m}}}.$$

By Proposition 2.26 (v), we have $\bar{I}R_{\mathfrak{m}} = \overline{IR_{\mathfrak{m}}}$. The ideal \bar{I} is homogeneous by [HS06, Corollary 5.2.3], so $\bar{I}R_{\mathfrak{m}} \cap R = \bar{I}$. We have the following, from which the claim follows.

$$\bar{I} \subseteq \bigcap_{t>0} \left(\overline{IR_{\mathfrak{m}} + J^t R_{\mathfrak{m}}} \cap R \right) \subseteq \overline{IR_{\mathfrak{m}}} \cap R = \bar{I}.$$

□

We are now able to prove Theorem 5.1 in the case that k is algebraically closed.

Proof. If $\text{ht}(I) \geq l+1$, then $c(I) \geq E_{l+1}(I) > E_l(I)$, so we must have $\text{ht}(I) = l$. Let $\ell_{l+1}, \dots, \ell_n$ be general linear forms and set $L = V(\ell_{l+1}, \dots, \ell_n)$. By Proposition 2.42, we have $E_l(I) = E_l(I|_L) \leq c(I|_L) \leq c(I)$, so we again have $E_l(I) = c(I|_L)$. We'll compute $d_1(I|_L), \dots, d_l(I|_L)$. Field extensions are faithfully flat, so $\text{ht}([I]_{\leq t}|_L)$ is preserved under field extension, hence so are the quantities $d_1(I|_L), \dots, d_l(I|_L)$. We may without loss of generality assume k is uncountable and algebraically closed.

Let $\gamma \in \text{GL}_n(k)$ be as in Lemmas 5.5 and 5.10 with respect to the ideal $I|_L$, and set $\mathfrak{a}_m = \text{in}_{>}(\gamma(I|_L)^m) = \text{gin}_{>}((I|_L)^m)$. As $c(I|_L) = E_l(I)$, by Corollary 3.11 we have

$$(36) \quad \overline{\mathbb{R}_{\geq 0}^l \setminus \Gamma(\mathfrak{a}_{\bullet})} = \text{conv} \left(\mathbf{0}, e_1(I|_L) \mathbf{b}_1^{(l)}, \frac{e_2(I|_L)}{e_1(I|_L)} \mathbf{b}_2^{(l)}, \dots, \frac{e_l(I|_L)}{e_{l-1}(I|_L)} \mathbf{b}_l^{(l)} \right).$$

The quantities $p_n(1), \dots, p_l(l)$ can be read off from Equation (36), so we have $\frac{e_j(I|_L)}{e_{j-1}(I|_L)} = d_j(I|_L)$ for all $1 \leq j \leq l$.

For $1 \leq j \leq l$, let f_j be an element of $[I]_{d_j}$ whose image mod $(\ell_{l+1}, \dots, \ell_n)$ is general. As $\text{ht}[I|_L]_{d_j} \geq j$, it follows that $f_1, \dots, f_l, \ell_{l+1}, \dots, \ell_n$ is a regular sequence. Let $J = (f_1, \dots, f_l)$. By Lemma 3.8 we have $E_l(J) = E_l(I)$. For $t > 0$, set $J_t = J + (\ell_{l+1}^t, \dots, \ell_n^t)$ and set $I_t = I + (\ell_{l+1}^t, \dots, \ell_n^t)$. Since $J_t \subseteq I_t$ is a complete intersection, Lemma 3.8 and Propositions 2.10 and 2.39 give

$$(37) \quad E_l(J) + \frac{n-l}{t} = E_l(J_t) \leq E_l(I_t).$$

Combining Equation (37) with Theorem 3.9 and Proposition 2.10 (v) gives

$$E_l(J) + \frac{n-l}{t} \leq E_l(I_t) \leq c(I_t) \leq c(I) + c((\ell_{l+1}^t, \dots, \ell_n^t)) = E_l(I) + \frac{n-l}{t}.$$

In particular, we have $E_l(I_t) = E_l(J_t) = c(J_t)$. Set

$$\mathfrak{D}_t := \overline{(x_1^{d_1(I)}, \dots, x_l^{d_l(I)}, x_{l+1}^t, \dots, x_n^t)}.$$

By Proposition 4.30, there exists $\gamma \in \mathrm{GL}_n(k)$ such that $\gamma_t \overline{J_t} = \mathfrak{D}_t$. Consequently, as $\mathfrak{D}_t \subseteq \gamma_t \overline{I_t}$ and $E_l(I_t) = E_l(\mathfrak{D}_t)$, by Proposition 2.39 we have $\overline{\mathfrak{D}_t} = \overline{I_t}$.

Set $\mathfrak{D} = \overline{(x_1^{d_1(I)}, \dots, x_l^{d_l(I)})}$. Let d be the maximum degree of an irredundant generator of \mathfrak{D} or \overline{I} . The k -vector space of forms of degree $\leq d$ in R is finite-dimensional, so the infinite descending chain $[\mathfrak{D}_1]_{\leq d} \supseteq [\mathfrak{D}_2]_{\leq d} \supseteq \dots$ — the limit of which is $[\mathfrak{D}]_{\leq d}$ by Lemma 5.11 — must stabilize; there exists $t_0 \gg 0$ such that $[\mathfrak{D}_t]_{\leq d} = [\mathfrak{D}]_{\leq d}$ for all $t \geq t_0$. Similarly, there exists $t_1 \gg 0$ such that $[\overline{I_t}]_{\leq d} = [\overline{I}]_{\leq d}$ for all $t \geq t_1$. For $t = \max(t_1, t_2)$, we have

$$\gamma_t [\overline{I}]_{\leq d} = \gamma_t [\overline{I_t}]_{\leq d} = [\mathfrak{D}_t]_{\leq d} = [\mathfrak{D}]_{\leq d}.$$

As $\gamma_t \overline{I}, \mathfrak{D}$ are generated in degree $\leq d$, it follows that $\gamma_t \overline{I} = \mathfrak{D}$. \square

5.2. Perfect fields. The following lemmas, together with the algebraically closed case of Theorem 5.1, allow us to generalize to the case of a perfect field.

Lemma 5.12. *Let L be a field. For $1 \leq i \leq r$, let $\mathbf{x}_i = x_{i,1}, \dots, x_{i,a_i}$ be a tuple of variables over L . Set $S = L[\mathbf{x}_1, \dots, \mathbf{x}_r, y_1, \dots, y_s]$. Let $e_1 < \dots < e_r$ be positive integers and set*

$$\mathfrak{D} = \overline{(x_{1,1}^{e_1}, \dots, x_{r,a_r}^{e_r})}.$$

If $\gamma \in \mathrm{GL}_{a_1 + \dots + a_r + s}$ fixes the flag

$$\left(0 \subseteq \mathrm{span}_L(\mathbf{x}_1) \subseteq \dots \subseteq \mathrm{span}_L(\mathbf{x}_1, \dots, \mathbf{x}_r) \right),$$

then $\gamma \mathfrak{D} = \mathfrak{D}$.

Proof. Let $w_{\mathbf{e}}$ denote the monomial valuation with $w_{\mathbf{e}}(\mathbf{x}_{i,j}) = \frac{1}{e_i}$ and $w_{\mathbf{e}}(y_i) = 0$. Then \mathfrak{D} consists precisely of the elements $f \in S$ such that $w_{\mathbf{e}}(f) \geq 1$. By the assumption on γ , we have $\gamma(x_{i,j}^{e_i}) \in (\mathbf{x}_1, \dots, \mathbf{x}_i)^{e_i}$, hence $w_{\mathbf{e}}(\gamma x_{i,j}^{e_i}) \geq 1$. It follows that

$$\gamma \mathfrak{D} = \overline{(\gamma x_{1,1}^{e_1}, \dots, \gamma x_{r,a_r}^{e_r})} \subseteq \overline{\mathfrak{D}} = \mathfrak{D}.$$

By symmetry, we conclude that $\gamma \mathfrak{D} = \mathfrak{D}$. \square

Lemma 5.13. *Let k be a field and L/k a separable field extension. Set $R = k[x_1, \dots, x_n]$ and $S = L[x_1, \dots, x_n]$. Let $d_1, \dots, d_l \in \mathbb{Z}^+$ and set $\mathfrak{D} = \overline{(x_1^{d_1}, \dots, x_l^{d_l})} \subseteq R$. Suppose that $I \subseteq R$ is a homogeneous ideal and that there exists $\gamma_L \in \mathrm{GL}_n(L)$ such that*

$$(38) \quad \gamma_L \overline{I S} = \mathfrak{D} S.$$

Then there exists $\gamma_k \in \mathrm{GL}_n(k)$ such that

$$(39) \quad \gamma_k \overline{I} = \mathfrak{D}.$$

Proof. By [HS06, Propositions 19.1.1, 19.1.2 and Corollary 19.5.2], we have $\overline{I S} = \overline{I S}$, so we may replace I by \overline{I} and ignore the integral closure in Equations (38) and (39).

Suppose that d_1, \dots, d_l take the values e_1, \dots, e_r with multiplicities a_1, \dots, a_r respectively and that $e_1 < \dots < e_r$. Re-index the variables $x_1, \dots, x_l, x_{l+1}, \dots, x_n$ as $\mathbf{x}_1, \dots, \mathbf{x}_r, y_{l+1}, \dots, y_n$ where $\mathbf{x}_i = x_{i,1}, \dots, x_{i,a_i}$. For $1 \leq i \leq r$, let I_i denote

the ideal generated by $[I]_{\leq e_i}$. Since L/k is separable, the ring $S/\sqrt{I_i}S$ is reduced, so $\sqrt{I_i}S = \sqrt{I_i}\overline{S}$. As $\sqrt{I_i}\overline{S} = \gamma_L^{-1}(\mathbf{x}_1, \dots, \mathbf{x}_i)$, it follows that $\sqrt{I_i}$ is generated by $a_1 + \dots + a_i$ linear forms.

Let $\gamma_k \in \mathrm{GL}_n(k)$ such that

$$\gamma_k \left(\left[\sqrt{I_1} \right]_1 \subseteq \dots \subseteq \left[\sqrt{I_r} \right]_1 \right) = \left(\mathrm{span}_k(\mathbf{x}_1) \subseteq \dots \subseteq \mathrm{span}_k(\mathbf{x}_1, \dots, \mathbf{x}_r) \right)$$

as flags in $\mathrm{span}(\mathbf{x}_1, \dots, \mathbf{x}_r)$. By construction, $\gamma_k \gamma_L^{-1} \in \mathrm{GL}_n(L)$ fixes the flag

$$\left(\mathrm{span}_L(\mathbf{x}_1) \subseteq \dots \subseteq \mathrm{span}_L(\mathbf{x}_1, \dots, \mathbf{x}_r) \right).$$

By Lemma 5.12, we deduce that

$$\gamma_k \overline{IS} = \gamma_k \gamma_L^{-1} \gamma_L \overline{IS} = \gamma_k \gamma_L^{-1} \mathfrak{D}S = \mathfrak{D}S.$$

By faithful flatness of $R \rightarrow S$, we conclude

$$\gamma_k I = (\gamma_k I)S \cap R = (\mathfrak{D}S) \cap R = \mathfrak{D}.$$

□

We now deduce Theorem 5.1.

Proof of Theorem 5.1. Set $L = \overline{k}$ and let $S = L[x_1, \dots, x_n]$. By Proposition 2.10 (vi) and Lemma 2.43 we have $c(IS) = c(I) = E_l(I) = E_l(IS)$. By Theorem 5.1 in the algebraically closed case, it follows that there exists $\gamma_L \in \mathrm{GL}_n(L)$, $d_1, \dots, d_l \in \mathbb{Z}^+$ such that Equation (38) holds. Because k is perfect, the extension L/k is separable, so Lemma 5.13 implies that there exists $\gamma_k \in \mathrm{GL}_n(k)$ such that Equation (39) holds, proving the theorem. □

To conclude this section, we note that the assumption on k cannot be weakened.

Example 5.14. Let k be a field of characteristic $p > 0$ and $t \in k \setminus k^p$. Let $R = k[x, y]$ and set $I = (x^p + ty^p)$. Then $c(I) = \frac{1}{p} = E_1(I)$, but I is not generated by a power of a linear form.

6. EXTENSIONS OF THEOREM B IN CHARACTERISTIC ZERO

One can ask whether Theorem 5.1 can be generalized to the local case. We note that the answer is “no” in positive characteristic, and we pose a conjecture generalizing Theorem 5.1 over the complex numbers.

Example 6.1. Let $R = \overline{\mathbb{F}}_p[[x, y]]$, and let $I = (x^p + y^{p+1}) \subseteq R$. Then $E_1(I) = \frac{1}{p} = \mathrm{fpt}(I)$, but $I = \overline{I}$ and there are no coordinates for R in which I is a monomial ideal.

For a complete description of when $E_1(I) = c(I)$ in positive characteristic, see [Bai26]. Whereas Theorem 5.1 fails to generalize to the local case in char $p > 0$, we are optimistic that a stronger result is possible in characteristic zero. Our conjecture extends [BA24, Remark 3.7] to the case of arbitrary codimension.

Conjecture 6.2. *Let $R = \mathbb{C}[[x_1, \dots, x_n]]$. Let $I \subseteq R$ be an ideal with $\text{ht}(I) \geq l$. Then $\text{lct}(I) = E_l(I)$ if and only if there exists a regular system of parameters y_1, \dots, y_n for R and positive integers d_1, \dots, d_l such that $\bar{I} = \overline{(y_1^{d_1}, \dots, y_l^{d_1})}$.*

We note that Conjecture 6.2, if proven, would be strong enough to give an alternate proof of Theorem 5.1 over \mathbb{C} .

Proof. Let $R = \mathbb{C}[[x_1, \dots, x_n]]$. Let I be a homogeneous ideal with $\text{ht}(I) \geq l$ and $E_l(I) = \text{lct}(I)$. Set $S = \mathbb{C}[[x_1, \dots, x_n]]$. Notation 2.9 and Definition 2.40 imply that $E_l(IS) = E_l(I)$ and $\text{lct}(IS) = \text{lct}(I)$, so by Conjecture 6.2 there exists a regular system of parameters y_1, \dots, y_n for S such that $\bar{IS} = \overline{(y_1^{d_1}, \dots, y_l^{d_1})}$.

Write each y_i as a power series in x_1, \dots, x_n and let z_1, \dots, z_n be the homogeneous degree-1 terms of y_1, \dots, y_n . As $y_i \equiv z_i \pmod{\mathfrak{m}^2}$, the forms z_1, \dots, z_n in R generate the homogeneous maximal ideal of R .

As I is homogeneous, so is \bar{I} , hence $\bar{I} = \bar{IS} \cap R$. In particular, we have $z_1^{d_1}, \dots, z_l^{d_l} \in \bar{I}$. By considering the homogeneous ideals $\bar{I} + \overline{(z_{l+1}^t, \dots, z_n^t)}$ for $t > 0$ (which evidently satisfy $E_n = \text{lct}$) and running an argument similar to Theorem 5.1, we deduce that $\bar{I} = \overline{(z_1^{d_1}, \dots, z_l^{d_l})}$. \square

We propose a stronger conjecture in terms of valuations. Let $R = \mathbb{C}[[x_1, \dots, x_n]]$. Let $v : R \rightarrow [0, \infty]$ be a valuation and let $A_R(v)$ be the log discrepancy as in [JM12]. For each positive integer n , define $\mathfrak{a}_n(v) = \{f \in R : v(f) \geq n\}$. By [JM12, Corollary 6.9] and [Blu18, Lemma 3.5], we have

$$(40) \quad \text{lct}(\mathfrak{a}_\bullet(v)) \leq A_R(v)$$

If v is centered on a prime ideal \mathfrak{p} of height at least l , then for $m \gg 0$ we have $\sqrt{\mathfrak{a}_m(v)} = \mathfrak{p}$ and so $\text{ht}(\mathfrak{a}_m(v)) \geq l$. Consequently, we have $E_l(\mathfrak{a}_\bullet(v)) \leq A_R(v)$.

Conjecture 6.3. *Let $R, v, \mathfrak{a}_n(v)$ be as above and assume v is centered on a prime ideal of height at least l . If $E_l(\mathfrak{a}_\bullet(v)) = A_R(v)$ then v is a monomial valuation.*

We conclude this article by demonstrating that Conjecture 6.3 implies Conjecture 6.2. First, a few preparatory results.

Definition 6.4. Let (R, \mathfrak{m}) be a Noetherian local ring and $I \subseteq R$ an ideal. The **fiber cone** of I is the ring

$$\mathcal{F}_I = \bigoplus_{n \geq 0} \frac{I^n}{\mathfrak{m}I^n}.$$

The **analytic spread** of I , denoted $\ell(I)$, is the dimension of \mathcal{F}_I .

Theorem 6.5 ([HS06], Theorem 8.3.7). *Let (R, \mathfrak{m}) be a Noetherian local ring such that R/\mathfrak{m} is infinite. If $I \subseteq R$ is a proper ideal, then $\ell(I)$ is equal to the least number of generators of an ideal $J \subseteq I$ such that $\bar{I} = \bar{J}$.*

Lemma 6.6. *Let (R, \mathfrak{m}) be a regular local ring of dimension n . Let $1 \leq l \leq n$ and let $I \subseteq J$ be ideals of height l . If $\sigma_l(I) = \sigma_l(J)$ and the analytic spread of J is equal to l , then $\bar{I} = \bar{J}$.*

Proof. Replacing R with $R[X]_{\mathfrak{m}_R[X]}$ changes neither the hypotheses nor the conclusion by [HS06, Lemma 8.4.2]; we may without loss of generality assume that R/\mathfrak{m} is infinite. By Proposition 2.37 (ii) and (iii), choose $f_1, \dots, f_l \in I$ such that, with $I' = (f_1, \dots, f_l)$, we have $\sigma_l(I) = \sigma_l(I')$. Moreover, by Theorem 6.5 we have $\ell(I') = l$.

Let $c_0(I), \dots, c_n(I)$ denote the **multiplicity sequence** of Achilles and Manaresi [AM97, Definition 2.2]. By [AMP19, Corollaries 2.6 and 2.7], for $K \in \{I', J\}$ we have $c_{n-l}(K) = \sigma_l(K)$. Moreover, by [AM97, Proposition 2.3 (i)] we have $c_i(K) = 0$ for $i \neq n - d$, so I', J have the same multiplicity sequence. It follows from [Pol+20, Theorem 4.2] that $\overline{I'} = \overline{J}$, which implies the claim. \square

Proposition 6.7. *If Conjecture 6.3 holds in dimension n and codimension l , then Conjecture 6.2 holds in dimension n and codimension l .*

Proof. Let $I \subseteq R$ be an ideal of height at least l such that $E_l(I) = \text{lct}(I)$. By Definition-Proposition 2.3, choose a divisorial valuation $v : R \rightarrow [0, \infty]$ such that $\text{lct}(I) = \frac{A_R(I)}{v(I)}$. Letting $w = \frac{v}{v(I)}$, we have $w(I) = 1$ and $A_R(w) = \text{lct}(I)$. As $w(I) = 1$, we have $I^n \subseteq \mathfrak{a}_n(w)$ for all $n > 0$, so $\text{ht}(w) \geq l$ and $E_l(\mathfrak{a}_\bullet(w)) \geq E_l(I)$.

It follows from Theorem 3.9 and eq. (40) that

$$(41) \quad \text{lct}(I) = E_l(I) \leq E_l(\mathfrak{a}_\bullet(w)) \leq \text{lct}(\mathfrak{a}_\bullet(w)) \leq A_R(w) = \text{lct}(I),$$

so in particular $E_l(\mathfrak{a}_\bullet(w)) = A_R(w)$. By Conjecture 6.3, w is a monomial valuation in some local coordinates z_1, \dots, z_n for R with $w(z_1) \geq w(z_2) \geq \dots \geq w(z_n)$. Since $\sigma_j(I) \leq \sigma_j(\mathfrak{a}_\bullet(w))$ for $1 \leq j \leq l$ and $E_l(I) = E_l(\mathfrak{a}_\bullet(w))$, we conclude $\sigma_j(I) = \sigma_j(\mathfrak{a}_\bullet(w))$ by [BA17, Proposition 10]. We can read off the invariants $\sigma_1(\mathfrak{a}_\bullet(w)), \dots, \sigma_l(\mathfrak{a}_\bullet(w))$ from the weights $w(z_i)$. In particular, we have $\sigma_j(\mathfrak{a}_\bullet(w)) = \frac{1}{w(z_1) \dots w(z_j)}$, hence $w(z_j) = \frac{\sigma_{j-1}(I)}{\sigma_j(I)}$ for all $1 \leq j \leq l$. Moreover, by definition of the log discrepancy for quasi-monomial valuations we have $A_R(w) = w(z_1) + \dots + w(z_n)$, so we must have $w(z_{l+1}) = \dots = w(z_n) = 0$.

By construction, $I \subseteq \mathfrak{a}_1(w)$. If the numbers $\frac{1}{w(z_1)}, \dots, \frac{1}{w(z_l)}$ are not all *integers*, then $\Gamma(\mathfrak{a}_1(w)) \subsetneq \Gamma(\mathfrak{a}_\bullet(w))$ – but this would imply

$$\text{lct}(I) \leq \text{lct}(\mathfrak{a}_1(w)) < \text{lct}(\mathfrak{a}_\bullet(w)),$$

which is impossible by Equation (41). Write $d_i = \frac{1}{w(z_i)}$ for $1 \leq i \leq l$. We are now in the situation where $I \subseteq \overline{(z_1^{d_1}, \dots, z_l^{d_l})} =: \mathfrak{D}$ and $E_l(I) = E_l(\mathfrak{D})$. Since \mathfrak{D} is the integral closure of an ideal generated by l elements, we have $\ell(\mathfrak{D}) = l$. By Lemma 6.6 we deduce that $\overline{I} = \overline{\mathfrak{D}}$. \square

The following argument is well-known to experts.

Proposition 6.8. *Conjecture 6.3 holds in codimension $l = 1$.*

Proof. If v is centered on a prime ideal of height 2 or more, then $\text{ht}(\mathfrak{a}_\bullet(v)) \geq 2$ and hence $A_R(v) \geq \text{lct}(\mathfrak{a}_\bullet(v)) \geq E_2(v) > E_1(v)$. Consequently, in order to have $E_1(v) = A_R(v)$, it must be the case that v is centered on a principal prime ideal

fR , hence $v = c \operatorname{ord}_f$ for some irreducible $f \in R$ and some $c \in \mathbb{R}^+$. We deduce that $E_1(f) = \operatorname{lt}(f)$, hence f is a power of a local coordinate by [GZ15]. \square

Proposition 6.9. *Conjecture 6.3 holds in dimension 2.*

Proof. By Proposition 6.8, it remains to consider the case $l = 2$. Suppose $E_2(v) = A_R(v)$. Without loss of generality, normalize v so that $v(\mathfrak{m}) = 1$. Let $\alpha(v) = \sup_{f \in \mathfrak{m}} v(f) / \operatorname{ord}_{\mathfrak{m}}(f)$ denote the *skewness* of α as in [FJ04]. By [BFJ14, Remark 4.9], we have $e_1(\mathfrak{a}_{\bullet}(v)) = e_2(\mathfrak{a}_{\bullet}(v)) = \frac{1}{\alpha(v)}$, so $E_2(v) = 1 + \alpha(v)$. By [FJ04, Proposition 3.48 (i)], we have $1 + \alpha(v) \leq A_R(v)$ with equality if and only if v is a monomial valuation, proving the claim. \square

REFERENCES

- [AM97] R. Achilles and M. Manaresi. “Multiplicities of a bigraded ring and intersection theory”. In: *Math. Ann.* 309.4 (1997), pp. 573–591. ISSN: 0025-5831,1432-1807. DOI: 10.1007/s002080050128.
- [AMP19] R. Achilles, M. Manaresi, and T. Pruschke. “Mixed multiplicities, Segre numbers and Segre classes”. In: *J. Algebra* 525 (2019), pp. 390–415. ISSN: 0021-8693,1090-266X. DOI: 10.1016/j.jalgebra.2019.01.028.
- [BA08] Carles Bivià-Ausina. “Joint reductions of monomial ideals and multiplicity of complex analytic maps”. In: *Math. Res. Lett.* 15.2 (2008), pp. 389–407. ISSN: 1073-2780. DOI: 10.4310/MRL.2008.v15.n2.a15.
- [BA17] Carles Bivià-Ausina. “Log canonical threshold and diagonal ideals”. In: *Proc. Amer. Math. Soc.* 145.5 (2017), pp. 1905–1916. ISSN: 0002-9939,1088-6826. DOI: 10.1090/proc/13382.
- [BA24] Carles Bivià-Ausina. *The log canonical threshold of products of ideals and mixed Łojasiewicz exponents*. 2024. arXiv: 2409.14251 [math.AC].
- [Bai25] Benjamin Baily. *On lower bounds for the F -pure threshold of equigenerated ideals*. 2025. arXiv: 2506.18891 [math.AC].
- [Bai26] Benjamin Baily. *Extremal F -thresholds in regular local rings*. 2026. arXiv: 2601.19885 [math.AC].
- [Ben+15] Angélica Benito et al. “Singularities of locally acyclic cluster algebras”. In: *Algebra Number Theory* 9.4 (2015), pp. 913–936. ISSN: 1937-0652,1944-7833. DOI: 10.2140/ant.2015.9.913.
- [BFJ14] Sébastien Boucksom, Charles Favre, and Mattias Jonsson. “A refinement of Izumi’s theorem”. In: *Valuation theory in interaction*. EMS Ser. Congr. Rep. Eur. Math. Soc., Zürich, 2014, pp. 55–81. ISBN: 978-3-03719-149-1.
- [Blu18] Harold Blum. “Existence of valuations with smallest normalized volume”. In: *Compos. Math.* 154.4 (2018), pp. 820–849. ISSN: 0010-437X,1570-5846. DOI: 10.1112/S0010437X17008016.
- [BS15] Bhargav Bhatt and Anurag K. Singh. “The F -pure threshold of a Calabi-Yau hypersurface”. In: *Math. Ann.* 362.1-2 (2015), pp. 551–567. ISSN: 0025-5831,1432-1807. DOI: 10.1007/s00208-014-1129-0.

- [DK01] Jean-Pierre Demailly and János Kollár. “Semi-continuity of complex singularity exponents and Kähler-Einstein metrics on Fano orbifolds”. In: *Ann. Sci. École Norm. Sup. (4)* 34.4 (2001), pp. 525–556. ISSN: 0012-9593. DOI: 10.1016/S0012-9593(01)01069-2.
- [DP14] Jean-Pierre Demailly and Hoang Hiep Pham. “A sharp lower bound for the log canonical threshold”. In: *Acta Math.* 212.1 (2014), pp. 1–9. ISSN: 0001-5962,1871-2509. DOI: 10.1007/s11511-014-0107-4.
- [DSB18] Alessandro De Stefani and Luis Núñez Betancourt. “ F -thresholds of graded rings”. In: *Nagoya Math. J.* 229 (2018), pp. 141–168. ISSN: 0027-7630,2152-6842. DOI: 10.1017/nmj.2016.65.
- [DSBP18] Alessandro De Stefani, Luis Núñez Betancourt, and Felipe Pérez. “On the existence of F -thresholds and related limits”. In: *Trans. Amer. Math. Soc.* 370.9 (2018), pp. 6629–6650. ISSN: 0002-9947,1088-6850. DOI: 10.1090/tran/7176.
- [Eis+] David Eisenbud et al. *ReesAlgebra: A Macaulay2 package. Version 2.3*. A *Macaulay2* package available at <https://github.com/Macaulay2/M2/tree/master/M2/Macaulay2/packages>.
- [Eis95] David Eisenbud. *Commutative algebra*. Vol. 150. Graduate Texts in Mathematics. With a view toward algebraic geometry. Springer-Verlag, New York, 1995, pp. xvi+785. ISBN: 0-387-94268-8; 0-387-94269-6. DOI: 10.1007/978-1-4612-5350-1.
- [EM21] Eva Elduque and Mircea Mustață. “On a conjecture of Teissier: the case of log canonical thresholds”. In: *Mat. Sb.* 212.3 (2021), pp. 175–192. ISSN: 0368-8666,2305-2783. DOI: 10.4213/sm9442.
- [FEM03] Tommaso de Fernex, Lawrence Ein, and Mircea Mustață. “Bounds for log canonical thresholds with applications to birational rigidity”. In: *Math. Res. Lett.* 10.2-3 (2003), pp. 219–236. ISSN: 1073-2780. DOI: 10.4310/MRL.2003.v10.n2.a9.
- [FEM04] Tommaso de Fernex, Lawrence Ein, and Mircea Mustață. “Multiplicities and log canonical threshold”. In: *J. Algebraic Geom.* 13.3 (2004), pp. 603–615. ISSN: 1056-3911,1534-7486. DOI: 10.1090/S1056-3911-04-00346-7.
- [Fer26] Tommaso de Fernex. Personal communication. 2026.
- [FJ04] Charles Favre and Mattias Jonsson. *The valuative tree*. Vol. 1853. Lecture Notes in Mathematics. Springer-Verlag, Berlin, 2004, pp. xiv+234. ISBN: 3-540-22984-1. DOI: 10.1007/b100262.
- [FM09] Tommaso de Fernex and Mircea Mustață. “Limits of log canonical thresholds”. In: *Ann. Sci. Éc. Norm. Supér. (4)* 42.3 (2009), pp. 491–515. ISSN: 0012-9593,1873-2151. DOI: 10.24033/asens.2100.
- [GS] Daniel R. Grayson and Michael E. Stillman. *Macaulay2, a software system for research in algebraic geometry*. Available at <http://www2.macaulay2.com>.

- [GW15] Jin Guo and Tongsuo Wu. “Monomial ideals under ideal operations”. In: *Comm. Algebra* 43.11 (2015), pp. 4745–4762. ISSN: 0092-7872,1532-4125. DOI: 10.1080/00927872.2014.952012.
- [GZ15] Qi’an Guan and Xiangyu Zhou. “Characterization of multiplier ideal sheaves with weights of Lelong number one”. In: *Adv. Math.* 285 (2015), pp. 1688–1705. ISSN: 0001-8708,1090-2082. DOI: 10.1016/j.aim.2015.08.002.
- [Her16] Daniel J. Hernández. “ F -purity versus log canonicity for polynomials”. In: *Nagoya Math. J.* 224.1 (2016), pp. 10–36. ISSN: 0027-7630,2152-6842. DOI: 10.1017/nmj.2016.14.
- [Hie25] Pham Hoang Hiep. *Singularity invariants of plurisubharmonic functions and complex spaces*. 2025. arXiv: 2304.02238 [math.CV].
- [How01] J. A. Howald. “Multiplier ideals of monomial ideals”. In: *Trans. Amer. Math. Soc.* 353.7 (2001), pp. 2665–2671. ISSN: 0002-9947,1088-6850. DOI: 10.1090/S0002-9947-01-02720-9.
- [HS06] Craig Huneke and Irena Swanson. *Integral closure of ideals, rings, and modules*. Vol. 336. London Mathematical Society Lecture Note Series. Cambridge University Press, Cambridge, 2006, pp. xiv+431. ISBN: 978-0-521-68860-4.
- [HT04] Nobuo Hara and Shunsuke Takagi. “On a generalization of test ideals”. In: *Nagoya Math. J.* 175 (2004), pp. 59–74. ISSN: 0027-7630,2152-6842. DOI: 10.1017/S0027763000008904.
- [Hun+08] Craig Huneke et al. “ F -thresholds, tight closure, integral closure, and multiplicity bounds”. In: vol. 57. Special volume in honor of Melvin Hochster. 2008, pp. 463–483. DOI: 10.1307/mmj/1220879419.
- [HY03] Nobuo Hara and Ken-Ichi Yoshida. “A generalization of tight closure and multiplier ideals”. In: *Trans. Amer. Math. Soc.* 355.8 (2003), pp. 3143–3174. ISSN: 0002-9947,1088-6850. DOI: 10.1090/S0002-9947-03-03285-9.
- [JM12] Mattias Jonsson and Mircea Mustață. “Valuations and asymptotic invariants for sequences of ideals”. In: *Ann. Inst. Fourier (Grenoble)* 62.6 (2012), pp. 2145–2209. ISSN: 0373-0956,1777-5310. DOI: 10.5802/aif.2746.
- [Kad+22] Zhibek Kadyrsizova et al. “Lower bounds on the F -pure threshold and extremal singularities”. In: *Trans. Amer. Math. Soc. Ser. B* 9 (2022), pp. 977–1005. ISSN: 2330-0000. DOI: 10.1090/btran/106.
- [Kim21] Dano Kim. “On a question of Teissier”. In: *Ann. Mat. Pura Appl. (4)* 200.5 (2021), pp. 2305–2311. ISSN: 0373-3114,1618-1891. DOI: 10.1007/s10231-021-01081-x.
- [Laz04] Robert Lazarsfeld. *Positivity in algebraic geometry. II*. Vol. 49. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]. Positivity

- for vector bundles, and multiplier ideals. Springer-Verlag, Berlin, 2004, pp. xviii+385. ISBN: 3-540-22534-X. DOI: 10.1007/978-3-642-18808-4.
- [LS81] Joseph Lipman and Avinash Sathaye. “Jacobian ideals and a theorem of Briançon-Skoda”. In: *Michigan Math. J.* 28.2 (1981), pp. 199–222. ISSN: 0026-2285,1945-2365.
- [May14] Sarah Mayes. “The limiting shape of the generic initial system of a complete intersection”. In: *Comm. Algebra* 42.5 (2014), pp. 2299–2310. ISSN: 0092-7872,1532-4125. DOI: 10.1080/00927872.2012.758271.
- [MTW05] Mircea Mustață, Shunsuke Takagi, and Kei-ichi Watanabe. “F-thresholds and Bernstein-Sato polynomials”. In: *European Congress of Mathematics*. Eur. Math. Soc., Zürich, 2005, pp. 341–364. ISBN: 3-03719-009-4.
- [Mus02] Mircea Mustață. “On multiplicities of graded sequences of ideals”. In: *J. Algebra* 256.1 (2002), pp. 229–249. ISSN: 0021-8693,1090-266X. DOI: 10.1016/S0021-8693(02)00112-6.
- [Mus12] Mircea Mustață. “IMPANGA lecture notes on log canonical thresholds”. In: *Contributions to algebraic geometry*. EMS Ser. Congr. Rep. Notes by Tomasz Szemberg. Eur. Math. Soc., Zürich, 2012, pp. 407–442. ISBN: 978-3-03719-114-9. DOI: 10.4171/114-1/16.
- [Pol+20] Claudia Polini et al. “Multiplicity sequence and integral dependence”. In: *Math. Ann.* 378.3-4 (2020), pp. 951–969. ISSN: 0025-5831,1432-1807. DOI: 10.1007/s00208-020-02059-5.
- [PRS24] Janet Page, Tim Ryan, and Karen E. Smith. *Smooth Surfaces with Maximal Lines*. 2024. arXiv: 2406.15868 [math.AG].
- [Ras15] Alexander Rashkovskii. “Extremal cases for the log canonical threshold”. In: *C. R. Math. Acad. Sci. Paris* 353.1 (2015), pp. 21–24. ISSN: 1631-073X,1778-3569. DOI: 10.1016/j.crma.2014.11.002.
- [Roc70] R. Tyrrell Rockafellar. *Convex analysis*. Vol. No. 28. Princeton Mathematical Series. Princeton University Press, Princeton, NJ, 1970, pp. xviii+451.
- [Sko72] Henri Skoda. “Sous-ensembles analytiques d’ordre fini ou infini dans \mathbf{C}^n ”. In: *Bull. Soc. Math. France* 100 (1972), pp. 353–408. ISSN: 0037-9484.
- [Smi20] Ilya Smirnov. “On semicontinuity of multiplicities in families”. In: *Doc. Math.* 25 (2020), pp. 381–400. ISSN: 1431-0635,1431-0643.
- [Tak04] Shunsuke Takagi. “F-singularities of pairs and inversion of adjunction of arbitrary codimension”. In: *Invent. Math.* 157.1 (2004), pp. 123–146. ISSN: 0020-9910,1432-1297. DOI: 10.1007/s00222-003-0350-3.
- [Tak06] Shunsuke Takagi. “Formulas for multiplier ideals on singular varieties”. In: *Amer. J. Math.* 128.6 (2006), pp. 1345–1362. ISSN: 0002-9327,1080-6377.
- [TW04] Shunsuke Takagi and Kei-ichi Watanabe. “On F-pure thresholds”. In: *J. Algebra* 282.1 (2004), pp. 278–297. ISSN: 0021-8693,1090-266X. DOI: 10.1016/j.jalgebra.2004.07.011.
- [TXZ24] Siyong Tao, Zida Xiao, and Huaiqing Zuo. *Bernstein-Sato roots for weighted homogeneous singularities in positive characteristic*. 2024. arXiv: 2410.20188v1 [math.AC].